

Dialkyl Carbonates as Sacrificial Molecules: from Heterocycles to Macrocycles

Fabio Aricò* and Pietro Tundo

Department of Environmental Science, Informatics and Statistics, Ca' Foscari University Venezia, Italy

*Corresponding author: fabio.arico@unive.it

Scientific topic: Catalysts, Pharma, Alternative solvents

Keywords: Cyclization, Heterocycles, Dialkyl carbonate, Green Chemistry, Chlorine-free Chemistry

Abstract

Dialkyl carbonate (DACs) are well recognized green reagents and solvents for new synthetic pathways. In particular dimethyl carbonate (DMC), nowadays synthesized by CO₂ insertion into epoxides, has shown surprising high selectivity with different nucleophiles acting either as methoxycarbonylation (B_{Ac}2 mechanism) or methylation (B_{Al}2 mechanism) agent.

In this lecture recent advances in DMC chemistry for chlorine-free synthesis of five- and six-membered heterocycles will be presented. Reaction of 1,4-diols with DMC in the presence of a base resulted in the chlorine-free synthesis of five-membered cyclic compounds. This synthetic procedure can be also used for the quantitative intramolecular heterocyclisation of bifunctional compounds, i.e., 4-amino-1-butanol to achieve pyrrolidine.^[1]




Figure 1. Synthesis of heterocycles by DMC chemistry

Six-member cyclic carbamates have also been synthesized by chlorine-free approach employing DMC chemistry. In fact, reacting a primary amine or a hydrazine with a di(methylcarbonate) derivative of 1,3-diols oxazinan-2-ones can be synthesized in a one-pot chlorine-free reaction.^[2]

Recently we also investigated the replacement of the chlorine by a carbonate moiety in half-nitrogen and -sulphur mustard compounds. Results collected demonstrated that the novel mustard carbonates are easily synthesized, don't show any toxicity and react with a wide range of nucleophiles in the absence of any base. These novel compounds can be employed for the synthesis of piperidines and thiopyrans. Furthermore, the polycondensation of a nitrogen mustard carbonate analogue with aromatic diols under dilution conditions resulted in a new synthetic approach to azacrowns previously not accessible.^[3]

References

- [1] F. Aricò, S. Evaristo, P. Tundo, "Synthesis of five- and six-membered heterocycles by dimethyl carbonate with catalytic amount of nitrogen bicyclic bases" *Green Chem.* 2015, 17:1176-1185.
- [2] C. R. McElroy, F. Aricò, P. Tundo, "1,3-Oxazinan-2-ones from amines and 1,3-diols through dialkyl carbonate chemistry" *Synlett* 2012, 23: 1809-1815
- [3] F. Aricò, I. Udrea, M. Crisma, P. Tundo, "Azacrown Ethers from Mustard Carbonate Analogues", *ChemPlusChem* 2015, 80:471-474.