

PAPER



Cite this: *Catal. Sci. Technol.*, 2016,
6, 6031

Received 12th February 2016,
Accepted 24th April 2016

DOI: 10.1039/c6cy00307a

www.rsc.org/catalysis

Hydration of aromatic alkynes catalyzed by a
self-assembled hexameric organic capsule†

Giorgio La Sorella,^a Laura Sporni,^a Pablo Ballester,^b Giorgio Strukul^a
and Alessandro Scarso^{*a}

The combination of a Brønsted acid catalyst and a supramolecular organic capsule formed by the self-assembly of six resorcin[4]arene units efficiently promotes the mild hydration of aromatic alkynes to their corresponding ketones. The capsule provides a suitable nanoenvironment that favors protonation of the substrate and addition of water.

Introduction

In recent years, supramolecular catalysis¹ is growing as a new cross-discipline that exploits the recognition properties of host structures to promote reaction catalysis through the implementation of weak intermolecular forces in the stabilization of reaction intermediates and transition states. The role of a catalytic host is to achieve high activity and both product and substrate selectivity² with the ultimate goal of enzyme mimicking.³ While unimolecular host structures featuring relatively small inner cavities, like cyclodextrins⁴ and cucurbiturils,⁵ have been investigated as catalysts in the past, more recently, supramolecular hosts displaying significantly larger cavities have become interesting. The increase in volume size of a container host enabled co-encapsulation of multiple substrates, as well as the nesting of metal catalysts and substrates. The inclusion of a metal complex catalyst can drastically alter its properties compared to those exhibited in bulk solution.⁶

As a function of solvent nature, different approaches were considered in the development of synthetic supramolecular catalysts. For example, water soluble capsular containers were assembled through the hydrophobic effect and explored as supramolecular catalysts for several chemical transformations in water.⁷ On the other hand, Rebek *et al.* introduced the assembly of hydrogen bonded capsules in organic sol-

vents as supramolecular catalysts for cycloaddition reactions.⁸

Resorcin[4]arene **1** is an easy to prepare molecule that, through hydrogen bonding interactions, spontaneously self-assembles⁹ into hexameric 1₆·(H₂O)₈, stable both in the solid state and in water saturated chloroform and benzene solutions. The hexameric capsule is characterized by a large cavity of about 1375 Å^{3,10} and encapsulates quaternary ammonium compounds¹¹ and some transition metal complexes by establishing cation-π interactions with the electron-rich internal lining of its aromatic rings.¹² Alternatively, the capsule interacts with hydrogen bonding species like alcohols¹³ or acids.¹⁴ These properties have been exploited to develop supramolecular catalytic systems based on the resorcin[4]arene hexameric capsule. In particular, the capsule has been used as (i) a reversible shield to control the activity of a photo-catalyst,¹⁵ (ii) a nano-reactor to impart unique substrate¹⁶ and product¹⁷ selectivities and (iii) a supramolecular catalyst itself.¹⁸

Some of the latter examples are related to the capsule mediated activation of water as a nucleophile towards the addition to different classes of substrates; in particular, the selective hydrolysis of acetals¹⁹ and the hydration of isonitriles to formamides driven by encapsulation of the substrates in the capsule that was recently disclosed by our group.²⁰

Alkyne hydration is an atom efficient reaction that transforms alkynes into carbonyl compounds such as ketones and aldehydes.²¹ The reaction is usually carried out with the aid of transition metal catalysts spanning from traditionally employed Hg species²² to more recent examples based on Au(I),²³ Pt(II)²⁴ and Ru(II).²⁵ Alternatively, the reaction can be catalyzed by strong Brønsted acids like sulphuric acid in extremely large excess,²⁶ as well as other strong organic acids that require harsh experimental conditions and long reaction times.²⁷ For example, intrinsically electron rich internal aryl alkynes require the presence of *p*-toluenesulfonic acid in

^a Dipartimento di Scienze Molecolari e Nanosistemi, Università Ca' Foscari di Venezia, via Torino 155, 30172, Mestre Venezia, ITALY. E-mail: alesca@unive.it; Fax: +39 041 2340517; Tel: +39 041 2348569

^b Institut for Research and Advanced Studies (ICREA) and Institut Català d'Investigació Química (ICIQ), The Barcelona Institute of Science and Technology, Avda. Països Catalans 16, 43007, Tarragona, Spain. E-mail: pballester@iciq.es; Fax: +34 977 920221


† Electronic supplementary information (ESI) available: Experimental details and GC-MS spectra. See DOI: 10.1039/c6cy00307a

ethanol solution under reflux conditions for several hours or heating with MW irradiation.²⁸ The use of microemulsion conditions in water with the aid of surfactants and 0.33 M HCl by heating the system from 100 to 140 °C for a few hours is also competent for the hydrolysis of alkynes.²⁹ Recently, it was also disclosed that the combination of sulphuric acid with an ionic liquid favoured the hydration of alkynes under mild reaction conditions.^{30,31} Likewise, the combination of a weak Brønsted acid as solvent with a strong Brønsted or Lewis acid catalyst like Ga(OTf)₃ enabled good yields of the carbonyl compound by heating at 100 °C for a few hours with reduced catalyst loading.³²

Results and discussion

Herein, we present an example of very efficient supramolecularly promoted hydration of terminal aromatic alkynes mediated by the combination of a strong Brønsted acid with the self-assembled supramolecular organic capsule of resorcin[4]arene 1₆·(H₂O)₈ (Scheme 1). The capsule acts as an organocatalyst and in the presence of a Brønsted acid efficiently transforms aromatic alkynes into their corresponding ketones at reaction temperatures of 60 °C or lower, within a few hours. The catalytic effect was observed exclusively when the capsule cavity was accessible. The addition of competitive guests that preferentially encapsulate inhibited the catalytic activity of the hexamer. This behaviour resembles the functioning of the active sites of enzymes.

The hydration reaction of phenylacetylene **2a** as model substrate was monitored over time by means of ¹H NMR spectroscopy. The reaction was initially investigated at 60 °C and for just a few hours. In the presence of the sole supramolecular capsule 1₆·(H₂O)₈, no reaction was observed (Table 1, entry 1). Similarly, in the presence of only strong Brønsted acids such as HBF₄, HCl or HNO₃, the hydrolysis reaction did not proceed (Table 1, entries 2–4). Only methanesulfonic acid, as expected,^{27a} and because of its better solubility in chloroform, led to the partial formation of acetophenone **3a**



Scheme 1 Hydration reaction of terminal aromatic alkynes **2** leading to their corresponding ketones **3** in the presence of tetrafluoroboric acid and mediated by the capsule 1₆·(H₂O)₈ and the competitive guest tetraethyl ammonium tetrafluoroborate **4**.

and small amounts of products derived from acid addition that were confirmed by GC–MS (Table 1, entries 5–6). Only when catalytic amounts of the capsule 1₆·(H₂O)₈ were combined with strong Brønsted acids did the quantitative hydration of **2a** to its corresponding acetophenone **3a** take place. In particular, using 10 mol% of the capsule and 50 mol% of HBF₄ with respect to the substrate **2a** catalyzed quantitative hydration in slightly more than one hour (Table 1, entry 7).

Analogous results were obtained using longer reaction times (18 hours) and by decreasing both the amount of the capsule and the acid (Table 1, entry 8). Similarly, HCl and methanesulfonic acid in combination with the capsule efficiently promoted the hydration reaction of **2a**. However, in this case, a decrease in the amounts of both species had a detrimental effect on the catalytic activity of the mixture (Table 1, entries 9–10 and 12–13). Conversely, the use of nitric acid as a Brønsted acid in the hydration reaction was not improved by the addition of 1₆·(H₂O)₈ (Table 1, entry 11).

It is widely accepted that the hydration reaction of alkynes occurs by water addition to a previously activated alkyne through protonation or coordination to a metal center. Both reaction mechanisms involve the enhancement of the electrophilicity of the substrate, thus favoring the addition of water. Recently, Tiefenbacher and co-workers demonstrated that the resorcin[4]arene hexamer behaves as a weak acid with a pK_a of about 5.5,¹⁹ while free resorcinol, not bound as in **1a**, has a pK_a of 9.15. The weak acidity of the capsule is not enough to induce the protonation of the substrate to a significant extent, as shown by the lack of reactivity in the presence of the capsule alone (Table 1 entry 1). The cooperation of the capsular hexamer with strong Brønsted acids like HBF₄ and HCl turned out to be much more efficient to catalyze the hydration reaction. As described above, it is well established that the hexamer favors the encapsulation of cationic guests and cationic reaction intermediates.^{18,33} Thus, a plausible, albeit not proven, explanation of the catalytic effect exerted by the capsule in the hydration of alkynes would possibly reside in the stabilization of positively charged species that are formed by Brønsted acid protonation of the triple bond.

In order to support this hypothesis, a series of control experiments were carried out. In order to exclude the involvement of the resorcinol moieties of **1** in the reaction, we replaced the hexameric capsule with 24 equivalents of resorcinol with respect to the original amount of capsule. We observed that under these reaction conditions, the formation of **3a** was negligible (Table 1, entry 16). We repeated the same control experiment using 4-*n*-hexyl-resorcinol as a more soluble resorcinol derivative observing a 39% yield of **3a** only after 24 h at 60 °C. These experiments were indicative of the minor effect exerted by the resorcinol units to the reaction catalysis.

The hydration of **2a** was also investigated in the presence of HBF₄, the capsule and tetraethylammonium tetrafluoroborate **4** as a competitive cationic guest for the capsule's interior.³⁴ The tetralkyl ammonium guest **4** was rapidly encapsulated as demonstrated by the appearance of a broad resonance at –0.05 ppm in the ¹H NMR spectrum (Fig. 1) with concomitant inactivation

Table 1 Catalytic tests for the hydration of **2a**

#	Brønsted Acid	Acid/2a	Time (h)	1 ₆ -8H ₂ O	4	3a (%) ^a
1	—	—	24	+	—	0
2	HBF ₄	0.5	1	—	—	<2
3	HCl	0.5	1	—	—	<2
4	HNO ₃	0.5	1	—	—	<2
5	CH ₃ SO ₃ H	0.5	1	—	—	57 (19) ^e
6 ^b	CH ₃ SO ₃ H	0.2	18	—	—	13 (3) ^e
7	HBF ₄	0.5	1.2	+	—	>98
8 ^b	HBF ₄	0.2	18	+	—	>98
9	HCl	0.5	1.5	+	—	>98
10 ^b	HCl	0.2	24	+	—	10
11	HNO ₃	0.5	1	+	—	18
12	CH ₃ SO ₃ H	0.5	1	+	—	>98
13 ^b	CH ₃ SO ₃ H	0.2	24	+	—	59 (17) ^e
14	HBF ₄	0.5	24	+	+	<2
15	HBF ₄	0.5	24	—	+	<2
16 ^c	HBF ₄	0.5	24	—	—	<2
17 ^d	HBF ₄	0.1	24	+	—	>98
18 ^f	HBF ₄	0.5	24	—	—	39

Experimental conditions: [**1**] = 36 mM, [**2a**] = 60 mM, [Brønsted acid] = 30 mM, [**4**] = 60 mM (10 eq. with respect to the capsule), water saturated chloroform-d = 1.5 mL, *T* = 60 °C. +: presence; -: absence. ^a Determined by ¹H-NMR. ^b [**1**] = 18 mM, [**2a**] = 60 mM, [Brønsted acid] = 12 mM. ^c [resorcinol] = 140 mM (24 eq. with respect to **1**₆-8H₂O). ^d [**1**] = 18 mM, [**2a**] = 120 mM, [Brønsted acid] = 12 mM. ^e Amount of Brønsted acid addition products. ^f [4-*n*-hexyl-resorcinol] = 140 mM (24 eq. with respect to **1**₆-8H₂O).

of the catalytic activity of the hexamer (Table 1, entry 14 and Fig. 1E).

All the reported data suggest that the cavity of the hexamer must be empty to allow the reaction to proceed. The ¹H-NMR analysis of a solution containing the capsule, the substrate phenylacetylene **2a** or **2a** and HBF₄ did not show evidence of encapsulated species (Fig. 1C). Most likely, the reaction requires initial, rate-determining, alkyne protonation by the strong Brønsted acid. The resulting cationic species is stabilized by encapsulation followed by a rapid attack of water that leads to the methyl ketone **3a**. Indeed, what matters

is not the stabilization of the cationic species itself but the effect on the transition state yielding the charged intermediate; hence, both the substrate and vinyl cation are expected to be within the cavity even if the former is not macroscopically revealed by NMR.

The scope of the hydration reaction was investigated by applying the catalytic system obtained by the combination of HBF₄ and **1**₆-8H₂O to a wide range of aromatic terminal alkynes, as summarized in Table 2. With the aim of investigating the turnover properties of the catalytic system under investigation, catalytic tests were performed with 50 mol% of HBF₄ and 10 mol% of **1**₆-8H₂O at 60 °C for 1 h and with 10 mol% of HBF₄ and 5 mol% of **1**₆-8H₂O at the same temperature for 24 h.

The hydration reaction turned out to be extremely sensitive to the electron density of the aromatic ring of the alkyne, providing higher yields for the more electron rich substrates. Furthermore, *N*-protonable 3- and 4-ethynyl pyridine derivatives did not produce their corresponding ketones in the presence of the capsule. On the other hand, extremely electron rich substrates bearing methoxy substituents, *i.e.* *p*-methoxyphenylacetylene **2g** and 4-methoxy-2-methyl-ethynylbenzene **2h**, can react at 60 °C both in the presence of the free hexamer and with the hexamer encapsulating the tetra-alkylammonium salt **4**. These results indicate that for these activated substrates, the Brønsted acidity of HBF₄ was enough to catalyze the hydration reaction (Table 2, entries 6 and 7). In these cases, the presence of the capsule either filled by the cationic species **4** or in its free form is substantially not influencing the reactivity of those activated substrates.

Substrates bearing alkyl substituents in the aromatic ring showed a clear improvement of the hydration reaction in the presence of the capsule, as reported in Table 2 (entries 1–5). It is noteworthy that a gradual decrease in the yield of the




Fig. 1 ¹H NMR spectra in water saturated chloroform-d: (A) phenylacetylene **2a** (60 mM); (B) phenylacetylene **2a** (60 mM) and **1**₆-8H₂O (6 mM) after 18 h at 60 °C; (C) phenylacetylene **2a** (60 mM) with HBF₄ (0.5 eq.) and **1**₆-8H₂O (6 mM); (D) phenylacetylene **2a** (60 mM) with HBF₄ (0.5 eq.) and **1**₆-8H₂O (6 mM) after 70 minutes at 60 °C; (E) phenylacetylene **2a** (60 mM) with HBF₄ (0.5 eq.), **1**₆-8H₂O (6 mM) and **4** (60 mM) after 24 h at 60 °C; ↓ encapsulated ammonium, ↑ acetophenone.

Table 2 Hydration reaction of **2b–2m** with HBF_4 mediated by the capsule $1_6 \cdot 8\text{H}_2\text{O}$























#	Substrate	Product	Yield (%) ^a
1			>98, 2 ^b , >98 ^c , 18 ^{b,c}
2			56, <2 ^b , 78 ^c , <2 ^{b,c}
3			78, <2 ^b , 40 ^c , <2 ^{b,c}
4			60, <2 ^b , 98 ^c , <2 ^{b,c}
5			47, 7 ^b , 83 ^c , 8 ^{b,c}
6			>98 ^c , 72 ^{b,c}
7			>98, >98 ^b
8			40, <2 ^b , 85 ^c , 3 ^{b,c}
9			54, 17 ^b

Table 2 (continued)

#	Substrate	Product	Yield (%) ^a
10			50 ^d , <2 ^{b,d}
11			25 ^d , <2 ^{b,d}

Experimental conditions: $[1] = 36 \text{ mM}$, $\text{HBF}_4/\text{substrate} = 0.5$, $[2\mathbf{b}-\mathbf{m}] = 60 \text{ mM}$, water saturated chloroform- $d = 1.5 \text{ mL}$, $T = 60 \text{ }^\circ\text{C}$, time = 1 h. ^a Determined by $^1\text{H NMR}$. ^b Reaction in the presence of $[4] = 60 \text{ mM}$. ^c $[1] = 18 \text{ mM}$; $\text{HBF}_4/\text{substrate} = 0.1$, $T = 60 \text{ }^\circ\text{C}$; time = 24 h. ^d $[1] = 180 \text{ mM}$; $\text{HBF}_4/\text{substrate} = 0.5$.

corresponding methyl ketone **3** was observed by increasing the size of the aromatic substrate **2**, regardless of its electronic nature (Table 2, entries 1–5). This suggests that the encapsulation of the protonated intermediate is size selective. The cavity of the hexamer can accommodate about 6–8 chloroform or benzene molecules. The encapsulation of the cationic intermediate of the reaction must occur together with a finite number of solvent molecules. In all cases, an agreement among the typical packing coefficient values observed for host-guest systems in solution is mandatory.³⁵ Further evidence of the selective encapsulation of the cationic intermediate was provided by the hydration reactions of large substrates such as 9-ethynyl-phenanthrene **2i** and 1-ethynyl-4-phenoxybenzene **2j** that led to much lower conversion to the corresponding ketone compared to the hydration of **2a** under similar experimental conditions (Table 2, entries 8 and 9). It is noteworthy that even with some differences related to the intrinsic reactivity of the different aromatic alkynes, the inhibition effect imparted by the presence of the competitive ammonium guest **4** was observed in almost all the substrates investigated, as shown in Table 2.

Other electron poor substrates like *p*-bromophenylacetylene, 1-(4-ethynylphenyl)ethanone and methyl-4-ethynylbenzoate failed to react in the presence of the capsule.

Finally, the hydration reaction was also tested using internal aromatic alkynes. We observed that 1-phenyl propyne **2l** and 1-phenyl-hexyne **2m** formed their corresponding ketones with moderate yields and required the presence of higher amounts of both the capsule and Brønsted acid (Table 2, entries 10 and 11). Conversely, 1,1'-ethyne-1,2-diylidibenzene and several terminal aliphatic alkynes did not show formation of their corresponding hydration products.³⁶

Conclusions

In conclusion, we have described another example of supramolecular catalysis in which catalytic amounts of the

hexameric self-assembled capsule $1_6\cdot 8\text{H}_2\text{O}$ in combination with sub-stoichiometric amounts of HBF_4 catalyzed the hydration of aromatic alkynes **2** to their corresponding methyl ketones **3** in a few hours using a water saturated chloroform solution heated at 60 °C. The Brønsted acid induces the protonation of the substrate, while the supramolecular capsule likely provides a suitable nano-environment to stabilize the transition state leading to the positively charged vinyl cation intermediate species,^{37,38} as would be expected with protonation being the rate determining step of the reaction. Moreover, we demonstrated the inactivation of the supramolecular catalyst by the addition of a competitive cationic ammonium guest **4** that is preferentially encapsulated. We also proved the sensitivity of the reaction to the size and shape of the substrates. These findings clearly speak for the occurrence of the hydration reaction within the cavity of the organocatalyst, thus mimicking the active site of enzymes.

Acknowledgements

AS acknowledges MIUR, Università Ca' Foscari di Venezia and Consorzio Interuniversitario Nazionale per la Scienza e Tecnologia dei Materiali for support. PB acknowledges the Gobierno de Espana MINECO for Project CTQ2014-56295-R, Severo Ochoa Excellence Accreditation (2014-2018 SEV-2013-0319); FEDER funds (Project CTQ2014-56295-R), and the ICIQ Foundation for funding.

Notes and references

- (a) M. Raynal, P. Ballester, A. Vidal-Ferran and P. W. N. M. van Leeuwen, *Chem. Soc. Rev.*, 2014, **43**, 1660; (b) M. Raynal, P. Ballester, A. Vidal-Ferran and P. W. N. M. van Leeuwen, *Chem. Soc. Rev.*, 2014, **43**, 1734.
- P. W. N. M. van Leeuwen, *Supramolecular Catalysis*, Wiley-VCH, 2008.
- (a) H. Wei and E. Wang, *Chem. Soc. Rev.*, 2013, **42**, 6060; (b) Y. Lin, J. Ren and X. Qu, *Acc. Chem. Res.*, 2014, **47**, 1097.
- R. Breslow and S. D. Dong, *Chem. Rev.*, 1998, **98**, 1997.
- K. I. Assaf and W. M. Nau, *Chem. Soc. Rev.*, 2015, **44**, 394.
- (a) P. Dydio and J. N. H. Reek, *Chem. Sci.*, 2014, **5**, 2135; (b) S. H. A. M. Leenders, R. Gramage-Doria, B. de Bruin and J. N. H. Reek, *Chem. Soc. Rev.*, 2015, **44**, 433; (c) J.-N. Rebilly, B. Colasson, O. Bistri, D. Over and O. Reinaud, *Chem. Soc. Rev.*, 2015, **44**, 467.
- J. H. Jordan and B. C. Gibb, *Chem. Soc. Rev.*, 2015, **44**, 547.
- (a) J. Kang and J. Rebek Jr, *Nature*, 1997, **385**, 50; (b) J. Kang, G. Hilmersson, J. Santamaria and J. Rebek, Jr., *J. Am. Chem. Soc.*, 1998, **120**, 3650; (c) J. Kang, J. Santamaria, G. Hilmersson and J. Rebek, Jr., *J. Am. Chem. Soc.*, 1998, **120**, 7389.
- L. R. MacGillivray and J. L. Atwood, *Nature*, 1997, **389**, 469.
- (a) L. Avram and Y. Cohen, *J. Am. Chem. Soc.*, 2002, **124**, 15148; (b) E. S. Barrett, T. J. Dale and J. Rebek Jr., *J. Am. Chem. Soc.*, 2008, **130**, 2344; (c) L. Avram and Y. Cohen, *Org. Lett.*, 2002, **4**, 4365; (d) L. Avram, Y. Cohen and J. Rebek, *Chem. Commun.*, 2011, **47**, 5368; (e) B. W. Purse and J. Rebek, *Proc. Natl. Acad. Sci. U. S. A.*, 2006, **103**, 2530; (f) A. Shivanyuk and J. Rebek, *J. Am. Chem. Soc.*, 2003, **125**, 3432; (g) M. Yamanaka, A. Shivanyuk and J. Rebek, *J. Am. Chem. Soc.*, 2004, **126**, 2939.
- A. Shivanyuk and J. Rebek Jr., *Proc. Natl. Acad. Sci. U. S. A.*, 2001, **98**, 7662.
- D. A. Dougherty, *Acc. Chem. Res.*, 2013, **46**, 885.
- (a) S. Slovak and Y. Cohen, *Chem. – Eur. J.*, 2012, **18**, 8515; (b) S. Slovak, L. Avram and Y. Cohen, *Angew. Chem., Int. Ed.*, 2010, **49**, 428; (c) B. Schnatwinkel, I. Stoll, A. Mix, M. V. Rekharsky, V. V. Borovkov, Y. Inoue and J. Mattay, *Chem. Commun.*, 2008, 3873; (d) O. Ugono and K. Travis Holman, *Chem. Commun.*, 2006, 2144.
- (a) Y. Aoyama, Y. Tanaka, Y. H. Toi and H. Ogoshi, *J. Am. Chem. Soc.*, 1988, **110**, 634; (b) Y. Aoyama, Y. Nonaka, Y. Tanaka, H. Toi and H. Ogoshi, *J. Chem. Soc., Perkin Trans. 1*, 1989, **2**, 1025.
- G. Bianchini, A. Scarso, G. La Sorella and G. Strukul, *Chem. Commun.*, 2012, **48**, 12082.
- (a) A. Cavarzan, J. N. H. Reek, F. Trentin, A. Scarso and G. Strukul, *Catal. Sci. Technol.*, 2013, **3**, 2898; (b) S. Giust, G. La Sorella, L. Sporni, G. Strukul and A. Scarso, *Chem. Commun.*, 2015, **51**, 1658.
- A. Cavarzan, A. Scarso, P. Sgarbossa, G. Strukul and J. N. H. Reek, *J. Am. Chem. Soc.*, 2011, **133**, 2848.
- (a) S. Shimizu, A. Usui, M. Sugai, Y. Suematsu, S. Shirakawa and H. Ichikawa, *Eur. J. Org. Chem.*, 2013, 4734; (b) G. La Sorella, L. Sporni, G. Strukul and A. Scarso, *ChemCatChem*, 2015, **7**, 291; (c) S. Giust, G. La Sorella, L. Sporni, F. Fabris, G. Strukul and A. Scarso, *Asian J. Org. Chem.*, 2015, **4**, 217; (d) L. Catti and K. Tiefenbacher, *Chem. Commun.*, 2015, **51**, 892; (e) Q. Zhang and K. Tiefenbacher, *Nat. Chem.*, 2015, **7**, 197.
- Q. Zhang and K. Tiefenbacher, *J. Am. Chem. Soc.*, 2013, **135**, 16213.
- G. Bianchini, G. La Sorella, N. Canever, A. Scarso and G. Strukul, *Chem. Commun.*, 2013, **49**, 5322.
- (a) L. Hintermann and A. Labonne, *Synthesis*, 2007, **8**, 1121; (b) L. Hintermann, *ChemCatChem*, 2012, **4**, 321; (c) L. Hintermann, *Top. Organomet. Chem.*, 2010, **31**, 123; (d) Y. Yamamoto, I. D. Gridnev, N. T. Patil and T. Jin, *Chem. Commun.*, 2009, 5075; (e) B. Xu, W. Wang, L.-P. Liu, J. Han, Z. Jin and G. B. Hammond, *J. Organomet. Chem.*, 2011, **696**, 269.
- (a) G. A. Olah and D. Meider, *Synthesis*, 1978, **9**, 671; (b) R. J. Thomas, K. N. Campbell and G. F. Hennion, *J. Am. Chem. Soc.*, 1938, **60**, 718.
- (a) W. E. Brenzovich Jr., *Angew. Chem., Int. Ed.*, 2012, **51**, 8933; (b) N. Marion, R. S. Ramón and S. P. Nolan, *J. Am. Chem. Soc.*, 2009, **131**, 448.
- F. Trentin, A. M. Chapman, A. Scarso, P. Sgarbossa, R. A. Michelin, G. Strukul and D. F. Wass, *Adv. Synth. Catal.*, 2012, **354**, 1095.
- (a) F. Boeck, T. Kribber, L. Xiao and L. Hintermann, *J. Am. Chem. Soc.*, 2011, **133**, 8138; (b) L. Li, M. Zeng and S. B. Herzon, *Angew. Chem., Int. Ed.*, 2014, **53**, 7892.

- 26 A. D. Allen, Y. Chiang, A. J. Kresge and T. T. Tidwell, *J. Org. Chem.*, 1982, **47**, 775.
- 27 (a) T. Tsuchimoto, T. Joya, E. Shirakawa and Y. Kawakami, *Synlett*, 2000, 1777; (b) N. Olivi, E. Thomas, J.-F. Peyrat, M. Alami and J.-D. Brion, *Synlett*, 2004, 2175; (c) N. Menashe and Y. Shvo, *J. Org. Chem.*, 1993, **58**, 7434.
- 28 M. Jacubert, O. Provot, J.-F. Peyrat, A. Hamze, J.-D. Brion and M. Alami, *Tetrahedron*, 2010, **66**, 3775.
- 29 Z. Nairoukh, D. Avnir and J. Blum, *ChemSusChem*, 2013, **6**, 430.
- 30 W.-L. Wong, K.-P. Ho, L. Y. S. Lee, K.-M. Lam, Z.-Y. Zhou, T. H. Chan and K.-Y. Wong, *ACS Catal.*, 2011, **1**, 116.
- 31 R. Kore and R. Srivastava, *Tetrahedron Lett.*, 2012, **53**, 3245.
- 32 S. Liang, G. B. Hammond and B. Xu, *Chem. Commun.*, 2015, **51**, 903.
- 33 L. Catti, Q. Zhang and K. Tiefenbacher, *Synthesis*, 2016, **48**, 313.
- 34 It is known that up to three molecules of tetraethylammonium tetrafluoroborate **4** are accommodated in the cavity of the resorcin[4]arene hexamer: L. C. Palmer, A. Shivanyuk, M. Yamanaka and J. Rebek Jr., *Chem. Commun.*, 2005, 857.
- 35 S. Mecozzi and J. Rebek Jr., *Chem. – Eur. J.*, 1998, **4**, 1016.
- 36 Diphenylacetylene and aliphatic alkynes like 4-phenyl-1-butyne, 3,3-dimethyl-1-butyne and 1-octyne did not lead to the formation of their corresponding ketones. Similarly, the hydration reaction of benzonitrile was tested under identical experimental conditions without observing the formation of the corresponding benzamide.
- 37 R. A. Cox, *Adv. Phys. Org. Chem.*, 2012, **46**, 1.
- 38 (a) A. D. Allen, Y. Chiang, A. J. Kresge and T. T. Tidwell, *J. Org. Chem.*, 1982, **47**, 775; (b) V. Lucchini and G. Modena, *J. Am. Chem. Soc.*, 1990, **112**, 6291.