



Ca' Foscari  
University  
of Venice

Department  
of Economics

Working Paper

Silvia Faggian  
Fausto Gozzi  
Peter M. Kort

Optimal investment with  
vintage capital:  
equilibrium distributions

ISSN: 1827-3580  
No. 12/WP/2019





## Optimal investment with vintage capital: equilibrium distributions

**Silvia Faggian**

*Ca' Foscari University of Venice*

**Fausto Gozzi**

*Università LUISS - Guido Carli*

**Peter M. Kort**

*CentER, Department of Econometrics & Operations Research, Tilburg University; Department of Economics, University of Antwerp*

### Abstract

The paper concerns the study of equilibrium points, or steady states, of economic systems arising in modelling optimal investment with vintage capital, namely, systems where all key variables (capitals, investments, prices) are indexed not only by time  $\tau$  but also by age  $s$ . Capital accumulation is hence described as a partial differential equation (briefly, PDE), and equilibrium points are in fact equilibrium distributions in the variable  $s$  of ages. Investments in frontier as well as non-frontier vintages are possible. Firstly a general method is developed to compute and study equilibrium points of a wide range of infinite dimensional, infinite horizon boundary control problems for linear PDEs with convex criterion, possibly applying to a wide variety of economic problems. Sufficient and necessary conditions for existence of equilibrium points are derived in this general context. In particular, for optimal investment with vintage capital, existence and uniqueness of a long run equilibrium distribution is proved for general concave revenues and convex investment costs, and analytic formulas are obtained for optimal controls and trajectories in the long run, definitely showing how effective the theoretical machinery of optimal control in infinite dimension is in computing explicitly equilibrium distributions, and suggesting that the same method can be applied in examples yielding the same abstract structure. To this extent, the results of this work constitutes a first crucial step towards a thorough understanding of the behaviour of optimal controls and trajectories in the long run.

### Keywords

Equilibrium Points, Equilibrium Distributions, Vintage Capital Stock, Age-structured systems, Maximum Principle in Hilbert Spaces, Boundary control, Optimal Investment

### JEL Codes

C61, C62, E22

*Address for correspondence:*

**Silvia Faggian**

Department of Economics  
Ca' Foscari University of Venice  
Cannaregio 873, Fondamenta S.Giobbe  
30121 Venezia - Italy  
Phone: (+39) 041 2346913  
e-mail: faggian@unive.it

*This Working Paper is published under the auspices of the Department of Economics of the Ca' Foscari University of Venice. Opinions expressed herein are those of the authors and not those of the Department. The Working Paper series is designed to divulge preliminary or incomplete work, circulated to favour discussion and comments. Citation of this paper should consider its provisional character.*

# OPTIMAL INVESTMENT WITH VINTAGE CAPITAL: EQUILIBRIUM DISTRIBUTIONS

SILVIA FAGGIAN [1], FAUSTO GOZZI [2], AND PETER M. KORT [3]

ABSTRACT. The paper concerns the study of equilibrium points, or steady states, of economic systems arising in modeling optimal investment with *vintage capital*, namely, systems where all key variables (capitals, investments, prices) are indexed not only by time  $\tau$  but also by age  $s$ . Capital accumulation is hence described as a partial differential equation (briefly, PDE), and equilibrium points are in fact equilibrium distributions in the variable  $s$  of ages. Investments in frontier as well as non-frontier vintages are possible. Firstly a general method is developed to compute and study equilibrium points of a wide range of infinite dimensional, infinite horizon boundary control problems for linear PDEs with convex criterion, possibly applying to a wide variety of economic problems. Sufficient and necessary conditions for existence of equilibrium points are derived in this general context. In particular, for optimal investment with vintage capital, existence and uniqueness of a long run equilibrium distribution is proved for general concave revenues and convex investment costs, and analytic formulas are obtained for optimal controls and trajectories in the long run, definitely showing how effective the theoretical machinery of optimal control in infinite dimension is in computing explicitly equilibrium distributions, and suggesting that the same method can be applied in examples yielding the same abstract structure. To this extent, the results of this work constitutes a first crucial step towards a thorough understanding of the behavior of optimal controls and trajectories in the long run.

*Key words:* Equilibrium Points; Equilibrium Distributions; Vintage Capital Stock; Age-structured systems; Maximum Principle in Hilbert spaces; Boundary control; Optimal Investment.

*Journal of Economic Literature:* C61, C62, E22

## 1. INTRODUCTION

Computing equilibrium points, or steady states, and describing their properties is one of the main goals in the mathematics of economic models. This task, when presuming an underlying optimal control problem with infinite horizon, is already nontrivial with one state variable, but it becomes harsh when the dynamics of the system are infinite dimensional, like in cases when heterogeneity/path dependency is taken into account. This is the case, for instance, of optimal investment with vintage capital (capital stock is heterogeneous in age, see *e.g.* [50, 35]), of spatial growth models (capital stock is heterogeneous in space, see *e.g.* [18, 34, 19]), of growth models with time-to-build (capital stock is path dependent, see *e.g.* [5, 6, 7]), or of models with heterogeneous agents (see

*e.g.* [67]). In all these examples, equilibrium points are indeed functions (of vintage, or space, or age) and may be more properly referred to as “equilibrium distributions”. Up to now such equilibrium distributions have been studied *only when the value function of the control problem is described by an analytic formula* – a requirement which is very seldom met – so that many interesting cases are left out of the picture.

On the contrary, this work addresses the study of equilibrium distributions in cases where no explicit formula for the value function is available, moreover it does so under the general assumptions of an infinite-horizon infinite-dimensional control problem with linear state equation and general convex (concave, in the application) payoff, providing a theoretical tool that can be used in a variety of applied examples. In fact, the theory is put immediately into practise for the optimal investment model with vintage capital, obtaining analytic formulas for the equilibrium distributions, and a complete sensitivity analysis for some instances of the problem. Hence the paper contains a theoretical and an applied part, both of equal weight and dignity, whose main achievements are listed below.

For the general theory (Sections 2, 3 and 4), we reprise and complete the study of the control problems analysed in Faggian and Gozzi [44].<sup>1</sup> There Dynamic Programming (DP) was employed to prove the existence and uniqueness of a regular solution  $v$  of the Hamilton-Jacobi-Bellman (HJB) equation, as well as a verification theorem implying existence and uniqueness of optimal feedback controls, and the fact that  $v$  coincides with the value function. Overall and differently from most contributions to the subject, this work presents an integrated approach between the DP and MP methods of optimal control theory. In particular:

- (a) a co-state is associated to the state variable, and necessary and sufficient conditions for the optimal path are established in the form of a Maximum Principle (MP) (Theorem 4.5);
- (b) the co-state associated to an optimal state is shown to coincide with the spatial gradient of the value function evaluated at that optimal state (Theorem 4.6);
- (c) the definition of two types of equilibrium points is introduced: the stationary solutions of the state-costate system, called MP-equilibrium points, and the stationary solutions of the closed loop equation (CLE) arising in the DP approach (Definition 4.8), called CLE-equilibrium points;
- (d) the relationship between the two types of equilibrium points is explained, and sufficient (and necessary) conditions for existence of such equilibria are provided (Theorem 4.10);
- (e) two results on the stability of CLE-equilibrium points are given by using or adapting the existing literature (Propositions 4.15 and 4.16).

---

<sup>1</sup>Note that in this theoretical context, since equilibrium distributions can be seen as points in a suitable infinite dimensional vector space, a space of functions, they will be still named “equilibrium points”

It is important noting that the theory cannot be used straightforwardly to treat applied problems in a satisfactory way. This happens on the one hand because the results in infinite dimension need to be translated into terms of the application under analysis, and on the other hand as it may be necessary to exploit the particular structure of the applied problem to specify formulas for practical use. One example is worked out through Theorem 5.5, in the case of the model of optimal investment with vintage capital.

In the applied part of this work (Sections 5 and 6), the theoretical results are used on the optimal investment model with vintage capital deriving:

- (e) the existence of MP- or CLE-equilibrium points, which is proven equivalent to the existence of solutions of a numerical equation explicitly derived from the data;
- (f) analytic formulas for MP- or CLE-equilibrium points in some relevant examples;
- (g) a sensitivity analysis for some particular sets of data.

In particular, the sensitivity analysis enables the development of new economic results while analyzing the vintage capital stock model in which revenue is a strictly concave and linear quadratic function of output, where the strict concavity is caused by market power on the output market. As is standard in this literature (Feichtinger et al. [50]), output linearly depends on the capital goods, whereas investment costs are convex and linear quadratic. We show that the equilibrium distribution capital stock is first increasing and then decreasing in the age of the capital good. The increasing part is the result of investment costs being relatively large when capital goods are relatively new. On the other hand, such investments are attractive due to the long lifetime of new capital goods. Capital goods of older age have a shorter lifetime. This gives an incentive to reduce investments in older capital goods, resulting in the fact that the equilibrium distribution capital stock for old machines decreases with respect to age. We further establish another non-monotonicity dependence of the equilibrium distribution capital goods level, but now with respect to the productivity of the capital goods. If productivity is relatively low, the number of capital goods increases if productivity goes up. This is because a given capital good produces more so that the firm is more eager to invest in it. On the other hand, if productivity is relatively large the firm decreases investments, because otherwise the firm overproduces resulting in a too low marginal revenue. In other words, some optimal output level exists and less capital goods are needed to produce this level when productivity is high.

In conclusion, this work shows how successfully and effectively the theoretical machinery of optimal control in infinite dimension is in computing *explicit* formulas and studying properties for equilibrium distributions, also *in absence of an explicit formula for the value function*. We believe that the theoretical tools developed in the first part of this work can be successfully employed in examples yielding the same abstract structure (like those mentioned at the beginning of this introduction) and possibly extended to more complex

cases with the use of suitable numerical approximations: this will be the subject of future work.

The paper is organized as follows. Section 2 presents a family of optimal investment models with vintage capital. Section 3 presents the abstract optimal control problem and shows that the problem contained in Section 2 falls into that wider class. Section 4 is the theoretical core of the paper, where we recall the results obtained with the DP approach in [44] (Section 4.1), we state and prove first order optimality conditions in terms of a Maximum Principle (Section 4.2), and we present and discuss the general results on equilibrium points (Section 4.3). In Section 5, the general results of the previous sections are applied to the model of optimal investment with vintage capital, providing a technique to derive analytic formulas for the equilibrium distributions. Finally, in Section 6, a sensitivity analysis is conducted on some instances of the problem of Section 5, i.e. where both revenues and costs are chosen linear-quadratic. This section also contains numerical results as illustration. An appendix with proofs of the theorems of Section 4 and 5, as well as some additional results, completes the work.

We remark that the paper is organized as to allow the reader less interested in mathematical details to approach Sections 5 and 6 without necessarily going through the theoretical Sections 3 and 4.

**1.1. Literature Review.** We complete this introductory section with an overview of literature on vintage capital, and on optimal control of infinite dynamical systems, thereby explaining what the present paper adds to each field.

From an economic point of view, the paper contributes to the literature of vintage capital stock models. Such models extend standard capital accumulation models, like, among many others, Eisner and Strotz [32] and Davidson and Harris [29] where capital goods are a function of just time. The extension is that also the age of the capital goods is taken into account. This enables to distinguish different vintages of capital goods so that one could explicitly analyze issues like aging (Barucci and Gozzi [14]), learning (Greenwood and Jovanovic [57]), pollution (Xepapadeas and De Zeeuw [72]), forest management (Fabbri, Faggian and Freni [37]), and technological progress (Feichtinger et al. [50]). We consider the kind of vintage capital stock models where investments in older capital goods are possible. This distinguishes the framework to be considered from works like Solow et al. [69], Malcomson [65], Benhabib and Rustichini [16], and Boucekkine et al. [20, 21, 22, 23].

The first contribution in vintage capital literature, which consider models where investments in older capital goods are also possible, is Barucci and Gozzi [15]. They consider the vintage capital stock framework where, as in Feichtinger et al. [49], revenue is linearly increasing in output, implying that the output price is constant, and linear-quadratic investment costs. Like in Feichtinger et al. [49], they do derive equilibrium distribution expressions for capital goods of different ages and corresponding investments. The present

paper generalizes these contributions by obtaining the equilibrium distribution expression of the capital goods for a model with general concave function.

Barucci and Gozzi [14] extends Barucci and Gozzi [15] by considering technological progress, while in Xepapadeas and De Zeeuw [72] the production process produces emissions next to products. Both papers keep the revenue linearly dependent on output. Provided an equilibrium distribution exists, which is not the case when we have ongoing technological progress as in Barucci and Gozzi [14], due to this linearity equilibrium distribution expressions are much easier to obtain compared to a revenue function being concave as in the present paper.

Closer to our present paper than the works cited above is Feichtinger et al. [50], in which also a firm with market power is considered. The difference with our work is that Feichtinger et al. considers technological progress. In particular, the main part of their work analyzes how the firm reacts with its investment policy to a technological breakthrough, which is a point in time at which a new technology is invented. The implication is that productivity of the capital goods of vintages borne after the breakthrough time jumps upwards. Our model is simpler in the sense that we do not consider technological progress. However, our analysis goes further than in Feichtinger et al. [50] in that we were able to derive an analytical expression for the equilibrium distribution. This we could do for a general concave revenue function, where Feichtinger et al. [50] just considers linear-quadratic revenue. Note that after the technological breakthrough Feichtinger's model turns into our model with prespecified revenue function. This implies that also in their framework a unique equilibrium distribution exists, which can be calculated using the results of the present paper.

From the point of view of mathematics, the main features of the optimal control problem here considered are: *(i)* the linear state equation and the convex cost criterion; *(ii)* the presence of a boundary control; *(iii)* the age structure of the driving operator  $A$  in the state equation.

Optimal control of infinite dimensional systems is the subject of many books and papers in the recent literature. Among the books in the deterministic case we mention Lions [62] and Barbu and Da Prato [9], and the more recent ones Li and Yong [63], and Troltzsch [71]. For the stochastic case (concerning the dynamic programming approach) one can see the recent book [36].

Concerning the dynamic programming approach to problems with linear state equation and convex cost but with distributed control, we refer the reader to Barbu and Da Prato [9, 10, 11], for some linear convex problems to Di Blasio [30, 31], for the case of constrained control to Cannarsa and Di Blasio [24], and for the case of state constraints to Barbu, Da Prato and Popa [12] (see also Gozzi [51, 52, 53] for a generalization of this approach to the case of semilinear state equations). For boundary control problems we recall, in the case of linear systems and quadratic costs (where the HJB equation reduces to the operator Riccati equation) *e.g.* the books by Lasiecka and Triggiani [60, 61], the book

by Bensoussan, Da Prato, Delfour and Mitter [17], and, for nonautonomous systems, the papers by Acquistapace, Flandoli and Terreni [1, 2, 3, 4]. For the case of a linear system and a general convex cost function, we mention the papers by Faggian [38, 39, 40, 41, 42], and by Faggian and Gozzi [43, 44] (in particular, the theory developed in the last two works is the starting point for theory in the present paper, and is recalled in Section 4.1). On the Pontryagin maximum principle for boundary control problems we mention again, in the linear quadratic case, the books [60, 61, 62] and [17]; in the case of linear systems with convex cost, e.g., the book by Barbu and Precupanu (Chapter 4 in [13]), and the papers [8], [58]; for general nonlinear boundary control problems, e.g., [27], [45], [46], [70] [55] [56]. None of them covers the class of problems treated here.

The main contributions of the present paper with respect to the mathematical literature quoted above are: (1) the proof of the Maximum Principle for infinite dimensional, infinite horizon optimal control problems with features (i) – (iii); (2) the co-state inclusion which reconnects the value function with the co-state; (3) the analysis of equilibrium points of the control problem.

## 2. THE OPTIMAL INVESTMENT MODEL WITH VINTAGE CAPITAL

We now describe the model of optimal investment with vintage capital, in the setting introduced by Barucci and Gozzi [15][14], and later reprised and generalized by Feichtinger et al. [48, 49, 50], and by Faggian [40, 41] and Faggian and Gozzi [43].

The capital accumulation process is given by the following system

$$(2.1) \quad \begin{cases} \frac{\partial K(\tau, s)}{\partial \tau} + \frac{\partial K(\tau, s)}{\partial s} + \mu K(\tau, s) = u_1(\tau, s), & (\tau, s) \in ]t, +\infty[ \times ]0, \bar{s}] \\ K(\tau, 0) = u_0(\tau), & \tau \in ]t, +\infty[ \\ K(t, s) = x(s), & s \in [0, \bar{s}] \end{cases}$$

with  $t > 0$  the initial time,  $\bar{s} \in [0, +\infty]$  the maximal allowed age, and  $\tau \in [0, T[$  with horizon  $T = +\infty$ . The unknown  $K(\tau, s)$  represents the amount of capital goods of age  $s$  accumulated at time  $\tau$ , the initial datum is a function  $x \in L^2(0, \bar{s})$  (the space of square integrable functions on  $(0, \bar{s})$ ),  $\mu > 0$  is a depreciation factor. Moreover,  $u_0 : [t, +\infty[ \rightarrow \mathbb{R}$  is the investment in new capital goods ( $u_0$  is the boundary control) while  $u_1 : [t, +\infty[ \times [0, \bar{s}] \rightarrow \mathbb{R}$  is the investment at time  $\tau$  in capital goods of age  $s$  (hence, the distributed control). Investments are jointly referred to as the control  $u = (u_0, u_1)$ . The output rate is

$$(2.2) \quad Q(K(\tau)) := \int_0^{\bar{s}} \alpha(s) K(\tau, s) ds,$$

where  $\alpha(s)$  is a productivity parameter. Selling the output to consumers results in an instantaneous revenue,  $R(Q)$ , where  $R$  is a concave function. Capital stock can be increased by investing, and investment costs are given by

$$(2.3) \quad C(u(\tau)) \equiv C_0(u_0(\tau)) + C_1(u_1(\tau)) \equiv C_0(u_0(\tau)) + \int_0^{\bar{s}} c_1(s, u_1(\tau, s)) ds,$$

with  $C_1$  indicating the investment cost rate for technologies of age  $s$ ,  $C_0$  the investment cost in new technologies, including adjustment costs,  $C_0, C_1$  convex in the control variables. The firm's payoff is then represented by the functional

$$(2.4) \quad I(t, x; u_0, u_1) = \int_t^{+\infty} e^{-\lambda\tau} [R(Q(K(\tau))) - C(u(\tau))] d\tau,$$

where  $\lambda \in \mathbb{R}$  is the discount rate. Note that  $\lambda$  is usually assumed positive, but here we leave the possibility of choosing a negative  $\lambda$  (corresponding, for example, to a negative interest rate). The entrepreneur's problem is that of maximizing  $I(t, x; u_0, u_1)$  over all state-control pairs  $\{K, (u_0, u_1)\}$  which are solutions (in a suitable sense) of equation (2.1) and keep the capital stock  $K(\tau, s)$  nonnegative at all times. Such a problem is known as *vintage capital* problem, for the capital goods depend jointly on time  $\tau$  and on age  $s$ , which is equivalent to their dependence on time and vintage  $\tau - s$ .

We finally recall the definition of the value function of the problem

$$(2.5) \quad V(t, x) := \inf_{u \in L^p_\lambda(t, +\infty, U)} I(t, x; u_0, u_1).$$

Since  $R$  and  $C$  are not time dependent it is immediate to see that

$$(2.6) \quad V(t, x) = e^{-\lambda t} V(0, x) =: V_0(x).$$

**Remark 2.1.** As a matter of fact, we treat the above problem without the state constraints  $K(\tau, s) \geq 0$  for all  $s$  and  $\tau$ , and check that constraints are satisfied *a posteriori* by the optimal trajectories of the unconstrained problem. In such a case, those trajectories are also optimal for the problem with state constraints. ■

**2.1. Revenues and costs.** In order to be able to treat optimal investment with vintage capital into the wider class of abstract problems described in Sections 3 and 4, we specify the assumptions on revenues  $R$  and costs  $C$  which ensure that the basic assumptions of the abstract problem (Assumptions 3.2, (3)-(6)) are fulfilled.

**Assumptions 2.2.** (i)  $R \in C^1(\mathbb{R})$ ,  $R$  concave,  $R'$  Lipschitz continuous. Moreover  $\alpha \in H^1(0, \bar{s})^2$  and  $\alpha(\bar{s}) = 0$ .  
(ii)  $C_0(r)$  and  $r \mapsto c_1(s, r)$  are convex, lower semi-continuous functions, with injective<sup>3</sup> subdifferential at all  $r \in \mathbb{R}$ .  
(iii)  $C_0^*(r), r \mapsto c_1(s, \cdot)^*(r)$  (are Fréchet differentiable and) have Lipschitz continuous derivatives, for all  $s \in [0, \bar{s}]$ .  
(iv)  $C_0(r)$  and  $r \mapsto c_1(s, r)$  are bounded below by a function of type  $a|r|^p + b$ , for some  $a > 0, b \in \mathbb{R}, p > 1$ .

<sup>2</sup> $H^1(0, \bar{s})$  is the space of square integrable functions which admit a square integrable derivative in weak sense. Continuous functions with piecewise continuous derivatives are included in this space.

<sup>3</sup>A multivalued function  $\rho : U \rightarrow \mathbb{R}$  is injective when  $\rho(u_1) \cap \rho(u_2) = \emptyset$  for every  $u_1, u_2 \in U, u_1 \neq u_2$ .

In the above statement, we denoted by  $f^*$  the convex conjugate of a convex function  $f$ , in particular  $C_0^*(r) = \sup_{w \in \mathbb{R}} \{wr - C_0(w)\}$ ,  $c_1(s, \cdot)^*(r) = \sup_{w \in \mathbb{R}} \{wr - c_1(s, w)\}$ . Note that no strong regularity of  $C$  is required.

For example, suitable choices for the *revenues* are the following:

- (a) *Linear-quadratic*:  $R(Q) = -aQ^2 + bQ$ ;
- (b) *Logarithmic*:  $R(Q) = \ln(1 + Q)$ , for  $Q \geq 0$  and  $R(Q) = Q$  for  $Q < 0$ ;
- (c) *Power*  $\gamma \in (0, 1)$ :  $R(Q) = b[(\nu + Q)^\gamma - \nu^\gamma]$ , with  $b, \nu > 0$  ( $\nu$  arbitrary small), for  $Q \geq 0$  and  $R(Q) = b\gamma\nu^{\gamma-1}Q$  for  $Q < 0$ . Note in particular that this  $R$  converges as  $\nu$  tends to 0 to  $R(Q) = bQ^\gamma$ , for  $Q \geq 0$  and  $R(Q) = -\infty$  for  $Q < 0$ <sup>4</sup>.

Suitable choices for the *costs* are, once set  $\beta = (\beta_0, \beta_1) \in \mathbb{R} \times L^\infty(0, \bar{s})$ , with  $\beta_1(s), \beta_0 \geq \epsilon \geq 0$ ,  $q = (q_0, q_1) \in \mathbb{R}_+ \times L^2(0, \bar{s})$ , the following:

(A) *Linear-quadratic*:

$$(2.7) \quad C(u) = \int_0^{\bar{s}} [\beta_1(s)u_1^2(s) + q_1(s)u_1(s)]ds + \beta_0u_0^2 + q_0u_0$$

(B) *Linear+quadratic with constrained control*:

$$(2.8) \quad C(u_0, u_1) = C_0(u_0) + C_1(u_1)$$

$$(2.9) \quad = q_0u_0 + g_{\beta_0, M_0}(u_0) + \int_0^{\bar{s}} [\alpha_1(s)u_1(s) + g_{\beta_1(s), M_1}(u_1(s))] ds$$

where

$$g_{\beta, M}(u) = \begin{cases} \beta u^2 & |u| \leq M \\ +\infty & |u| > M \end{cases}$$

Such a cost can be easily generalized to a case where  $u$  belongs to any compact interval and not necessarily  $u \in [-M, M]$ .

(C) *Linear+Power costs*:

$$(2.10) \quad C(u_0, u_1) = C_0(u_0) + C_1(u_1)$$

$$(2.11) \quad = q_0u_0 + f_{\beta_0}(u_0) + \int_0^{\bar{s}} [q_1(s)u_1(s) + f_{\beta_1(s)}(u_1(s))] ds$$

where, for  $p > 2$ ,

$$f_\beta(u) = \begin{cases} \beta [(u + \theta)^p - \theta^p] & u \geq 0 \\ +\infty & u < 0 \end{cases}$$

which implies also positivity constraints of the controls.

We treat all of these cases in Section 5. Moreover, in Section 6 we treat the case of linear–quadratic revenues and costs for which we derive analytic formulas for the long run optimal couples, and perform a complete sensitivity analysis.

---

<sup>4</sup>The definition of  $R$  for negative values of  $Q$  is needed in order to apply the general theory, although negative values of  $Q$  will never emerge in our calculations. Note also that setting  $R(Q) = -\infty$  for  $Q < 0$  is equivalent to require  $Q \geq 0$  in optimal solutions.

The reader is advised that Sections 3, 4, and the Appendix are devoted to the mathematics of the general problem and require a good knowledge of functional analysis to be fully understood. Nonetheless, they may be skipped at a first reading, as the reader will find in Section 5 the theoretical results translated in terms of the problem of optimal investment with vintage capital.

### 3. THE THEORETICAL FRAMEWORK

Here we introduce an abstract class of infinite dimensional optimal control problems with linear evolution equation and convex payoff, in which the control may also act on the boundary, and address it as (P). Then, in Section 3.3 we show that the optimal investment model with vintage capital described in the previous section is of type (P).

**3.1. Notation.** The expression  $a \vee b$  means the maximum of the real numbers  $a$  and  $b$ . If  $X$  is a Banach space, we indicate its norm with  $|\cdot|_X$ , its dual with  $X'$ , with  $\langle \cdot, \cdot \rangle_{X', X}$  the duality pairing. When  $X = V'$  we use for simplicity  $\langle \cdot, \cdot \rangle$  in place of  $\langle \cdot, \cdot \rangle_{V', V}$ . If  $X$  is also a Hilbert space, we indicate with  $(\cdot | \cdot)_X$  the inner product in  $X$ .

If  $X$  and  $Y$  are Banach spaces, then  $C^1(X)$  denotes all Fréchet differentiable functions from  $X$  to  $\mathbb{R}$ , and  $\mathcal{L}(X, Y)$  the set of all linear and continuous operators from  $X$  to  $Y$ , with associated norm  $\|\cdot\|_{\mathcal{L}(X, Y)}$ . Moreover we set

$$Lip(X; Y) = \{f : X \rightarrow Y : [f]_L := \sup_{x, y \in X, x \neq y} \frac{|f(x) - f(y)|_Y}{|x - y|_X} < +\infty\}$$

$$C_{Lip}^1(X) := \{f \in C^1(X) : [f']_L < +\infty\}$$

and, for  $p \geq 1$ ,

$$\mathcal{B}_p(X, Y) := \{f : X \rightarrow \mathbb{R} : |f|_{\mathcal{B}_p} := \sup_{x \in X} \frac{|f(x)|_Y}{1 + |x|_X^p} < +\infty\}, \quad \mathcal{B}_p(X) := \mathcal{B}_p(X, \mathbb{R}),$$

$$C([0, T], \mathcal{B}_p(X, Y)) := \{f : [0, T] \rightarrow \mathcal{B}_p(X, Y) : f \text{ continuous}\}$$

Note that  $\mathcal{B}_p$  are Banach spaces if endowed with the norm  $|\cdot|_{\mathcal{B}_p}$ , so that continuity is intended with respect to such norms. Furthermore, we set

$$\Sigma_0(X) := \{w \in C_{Lip}^1(X) : w \text{ is convex}\}.$$

Finally, if  $X$  is a Hilbert space and  $h : X \rightarrow \mathbb{R}$  is a convex function, then  $h^*$  will denote its convex conjugate, namely  $h^* : X \rightarrow \mathbb{R}$ ,  $h^*(x) = \sup_{y \in X} \{\langle x, y \rangle - h(y)\}$ .

**3.2. The abstract optimal control problem (P).** We consider two real separable Hilbert spaces  $V$  and  $H$  with  $V$  continuously embedded in  $H$ . We identify  $H$  with its dual and we call  $V'$  the topological dual of  $V$ , which we do not identify with  $V$  for the reasons explained in Section 3.3. We then get a so-called Gelfand triple

$$V \hookrightarrow H \hookrightarrow V'$$

We choose  $V'$  as state space. The control space is the real separable Hilbert space  $U$  (which we identify with its dual  $U'$ ). We consider the control system with state space  $V'$ , control space  $U$ , and varying initial time  $t \geq 0$ , described by

$$(3.1) \quad \begin{cases} y'(\tau) = Ay(\tau) + Bu(\tau), & \tau > t \\ y(t) = x \in V', \end{cases}$$

where  $A$  and  $B$  are linear operators, possibly unbounded. Moreover, we take a convex functional of the following type

$$(3.2) \quad J(t, x, u) = \int_t^{+\infty} e^{-\lambda\tau} [g_0(y(\tau)) + h_0(u(\tau))] d\tau$$

where the function  $g_0$  and  $h_0$  are convex functions. The problem (P) is that of minimizing  $J_\infty(t, x, u)$  with respect to  $u$ , over the set of admissible controls

$$(3.3) \quad L_\lambda^p(t, +\infty; U) = \{u : [t, +\infty) \rightarrow U ; \tau \mapsto u(\tau)e^{-\frac{\lambda\tau}{p}} \in L^p(t, +\infty; U)\},$$

which is a Banach space with the norm

$$\|u\|_{L_\lambda^p(t, +\infty; U)} = \left( \int_t^{+\infty} |u(\tau)|_U^p e^{-\lambda\tau} d\tau \right)^{\frac{1}{p}} = \|e^{-\frac{\lambda(\cdot)}{p}(\cdot)} u(\cdot)\|_{L^p(t, +\infty; U)}.$$

**Remark 3.1.** In the above problem no constraints on controls or on states are assumed although, in economic applications, the state represents capital stock, usually assumed nonnegative. Here we proceed along with the frequently used idea (see e.g.[35]) to check *ex post* that the constraints are satisfied by the optimal trajectories of the unconstrained problem, so those trajectories are optimal also for the constrained problem. ■

The basic assumptions on the data are stated below and will hold throughout the paper.

**Assumptions 3.2.** (1)  $A : D(A) \subset V' \rightarrow V'$  is the infinitesimal generator of a strongly continuous semigroup  $\{e^{\tau A}\}_{\tau \geq 0}$  on  $V'$ . Moreover there exists  $\omega \in \mathbb{R}$  such that<sup>5</sup>

$$\|e^{\tau A}x\|_{V'} \leq e^{\omega\tau} \|x\|_{V'}, \quad \forall \tau \geq 0;$$

$$(2) B \in L(U, V');$$

$$(3) g_0 \in \Sigma_0(V')$$

$$(4) h_0 : U \rightarrow \mathbb{R} \text{ is convex and lower semi-continuous, } \partial h_0 \text{ is injective.}$$

$$(5) h_0^*(0) = 0, h_0^* \in \Sigma_0(U);$$

$$(6) \exists a > 0, \exists b \in \mathbb{R}, \exists p > 1 : h_0(u) \geq a|u|_U^p + b, \forall u \in U;$$

$$(7) \lambda > \omega.$$

In proving some results we will need to specify the assumption (7) above as follows.

**Assumptions 3.3.** In addition to Assumption 3.2, we require that either

<sup>5</sup>When  $\omega > 0$ , a semigroup  $S(t)$  with this property is usually called a *pseudo-contraction semigroup*, as  $e^{-\omega t}S(t)$  is a contraction semigroup with generator  $A - \omega I$ .

$$(1) \quad p > 2, \quad \lambda > (2\omega) \vee \omega$$

or

$$(2) \quad g_0 \in \mathcal{B}_1(V').$$

The adjoint of  $A$  in the inner product of  $V'$  is denoted by  $A^*$ , while the adjoint of  $A$  with respect to the duality  $\langle \cdot, \cdot \rangle$  in  $V, V'$  is the unbounded operator  $A_1^*$  on  $V$ ,  $A_1^* : D(A_1^*) \subset V \rightarrow V$ .

**Remark 3.4.** We recall that, if a function  $h : U \rightarrow \mathbb{R} \cup \{+\infty\}$  is lower semicontinuous and convex (and not identically  $+\infty$ ), then the subgradient  $\partial h$  is defined as  $\partial h(u) = \{u^* \in U : h(v) - h(u) \geq \langle u^*, v - u \rangle_U, \forall v \in U\}$ . Moreover if  $\partial h$  is injective then  $h^*$  is Fréchet differentiable with  $(h^*)'(u) = (\partial h)^{-1}(u)$  for all  $u \in U$ . ■

**3.3. Optimal investment with vintage capital is of type (P) .** We end the section by showing that the problem of optimal investment with vintage capital described in Section 2 falls in the general class (P) described above and refer interested readers to [41] for full detail.<sup>6</sup>

We at first formulate an intermediate abstract problem in  $H = L^2(0, \bar{s})$ , the space of square integrable functions of variable  $s$ , using the modified translation semigroup  $\{e^{A_0 t}\}_{t \geq 0}$  on  $H$ , namely the linear operators  $e^{A_0 t} : H \rightarrow H$  such that

$$[e^{A_0 t} f](s) = f(s - t) e^{-\mu t}, \quad \text{if } s \in [t, \bar{s}], \quad \text{and} \quad [e^{A_0 t} f](s) = 0 \text{ otherwise.}$$

If  $H^1(0, \bar{s}) = \{f \in L^2(0, \bar{s}) : f' \in L^2(0, \bar{s})\}$ , then the generator of  $\{e^{A_0 t}\}_{t \geq 0}$  is the operator  $A_0 : D(A_0) \subset H \rightarrow H$ , with

$$D(A_0) = \{f \in H^1(0, \bar{s}) : f(0) = 0\}, \quad A_0 f(s) = -f'(s) - \mu f(s).$$

The adjoint of  $A_0$  is then  $A_0^* : D(A_0^*) \rightarrow H$  with

$$D(A_0^*) = \{f \in H^1(0, \bar{s}) : f(\bar{s}) = 0\}, \quad [A_0^* f](s) = f'(s) - \mu f(s),$$

generating itself a modified translation semigroup  $\{e^{A_0^* t}\}_{t \geq 0}$  on  $H$ , given by

$$[e^{A_0^* t} f](s) = f(s + t) e^{-\mu t}, \quad \text{if } s \in [0, \bar{s} - t], \quad \text{and} \quad [e^{A_0^* t} f](s) = 0 \text{ otherwise.}$$

The control space is  $U = \mathbb{R} \times H$ , the control function is a couple

$$u \equiv (u_0, u_1) : [t, +\infty) \rightarrow \mathbb{R} \times H,$$

and the control operator is given by

$$Bu \equiv B(u_0, u_1) = u_1 + u_0 \delta_0, \quad \text{for all } (u_0, u_1) \in \mathbb{R} \times H,$$

<sup>6</sup>See also [33], [17] or [68] for the general theory of strongly continuous semigroups and evolution equations.

$\delta_0$  being the Dirac delta at the point 0. With this notation, the original state equation (2.1) can be written as

$$(3.4) \quad \begin{cases} K'(\tau) = A_0 K(\tau) + Bu(\tau), & \tau > t \\ K(t) = x, \end{cases}$$

Note that  $H$  and  $U$  are Hilbert spaces, and that  $B$  is unbounded, meaning that it is *not* a continuous operator from  $U$  to  $H$  (unless  $u_0 = 0$ , corresponding to identically null boundary control  $u_0(\tau)$ ), for the Dirac delta does not lie in  $H$ . Then (3.4) needs to be interpreted in a suitable way, for instance in an extended state space.

Then we generalize all previous notions to a wider space. We set  $V \equiv D(A_0^*)$ , and assume  $V'$  as state space of the abstract problem. Indeed by standard arguments (see e.g. [33, Section II.5]) – and in particular by replacing the scalar product in  $L^2$  with the duality pairing  $\langle \phi, \psi \rangle$  with  $\phi \in V'$ ,  $\psi \in V$  (coinciding with the inner product in  $L^2$  when  $\phi \in L^2$ ) – the semigroup  $\{e^{A_0 t}\}_{t \geq 0}$  can be extended to a strongly continuous semigroup  $\{e^{A t}\}_{t \geq 0}$  on  $V'$ , by setting

$$(3.5) \quad \langle e^{A t} \phi, f \rangle = \langle \phi, e^{A_0^* t} f \rangle \quad \text{for every } f \in V, \phi \in V',$$

The generator of  $\{e^{A t}\}_{t \geq 0}$  is the operator  $A : D(A) \subset V' \rightarrow V'$ , with  $D(A) = H$ . Moreover the semigroup  $\{e^{A_0^* t}\}_{t \geq 0}$  can be restricted to a strongly continuous semigroup on  $V$ , with generator the restriction of  $A_0^*$  to  $D((A_0^*)^2)$ . Such restriction is exactly the adjoint of  $A$  in the duality  $\langle \cdot, \cdot \rangle$  and is then denoted, as in the previous subsection, by  $A_1^*$ .

The role of  $H$  is that of pivot space between  $V$  and  $V'$ , namely  $V \subset H \subset V'$ , with continuous inclusions. The control operator  $B$  is then in  $L(U, V')$ . Its adjoint is given by

$$(3.6) \quad B^* : V \rightarrow U, \quad \text{with } B^* v = (v(0), v).$$

It is also useful to note that  $A^{-1}$  is well defined and that

$$(3.7) \quad [-A^{-1} \delta_0](s) = e^{\mu s}, \quad \text{while } [-A^{-1} f](s) = \int_0^s e^{-\mu(s-\sigma)} f(\sigma) d\sigma, \quad \text{for all } f \in H$$

The target functional is also interpreted on extended spaces once the production function is described as

$$(3.8) \quad Q(K(\tau)) = \langle \alpha, K(\tau) \rangle$$

where, the duality pairing  $\langle \cdot, \cdot \rangle$  between  $V$  and  $V'$  has replaced the scalar product in  $L^2$  in the original definition (2.2) of  $Q$ , and  $c : U \rightarrow \mathcal{U}$ . Then (2.4) becomes

$$I(t, x; u_0, u_1) = \int_t^{+\infty} e^{-\lambda \tau} [R(\langle \alpha, K(\tau) \rangle) - C(u_0(\tau), u_1(\tau))] d\tau$$

The firm's optimal investment problem falls into the wider class described in the next theoretical sections, provided it is reformulated as a minimization problem, where the functions  $g_0$  and  $h_0$  there described are chosen as

$$(3.9) \quad g_0(x) := -R(\langle \alpha, x \rangle), \quad h_0(u_0, u_1) := C(u_0, u_1).$$

Indeed the following Lemma holds true.

**Lemma 3.5.** *Assumptions 2.2 imply, along with the above definitions of  $A$  and  $B$  and (3.9), that Assumptions 3.2 are satisfied with  $\omega = -\mu$ . Furthermore if  $p > 2$ , then Assumption 3.3 (1) is satisfied. If instead if  $p > 1$  and  $R$  has at most linear growth, Assumption 3.3 (2) is satisfied.*

*Proof.* Assumption 3.2-(1) is satisfied with  $\omega = -\mu$  since, for every  $\phi \in V'$  we have, by definition of  $V'$

$$|e^{\tau A}\phi|_{V'} = \sup_{|f|_V=1} \langle e^{\tau A}\phi, f \rangle = \sup_{|f|_V=1} \langle \phi, e^{\tau A_0^*} f \rangle \leq \sup_{|f|_V=1} |\phi|_{V'} |e^{\tau A_0^*} f|_V \leq e^{-\mu t} |\phi|_{V'}.$$

Assumption 3.2-(2) is trivially satisfied as pointed out above in the definition of  $B$ . By Assumption 2.2,  $\alpha \in V$  and  $g_0$  is a Fréchet differentiable convex function of  $x \in V'$ , with Fréchet differential  $g'_0(x)$  defined by  $g'_0(x)[s] = -R'(\langle \alpha, x \rangle)\alpha(s)$ . Such differential is a Lipschitz continuous function of  $x$ , with Lipschitz constant  $Lip(g'_0) = Lip(R')|\alpha|_V^2$ , as

$$(3.10) \quad |g'_0(x) - g'_0(y)|_V \leq |R'(\langle \alpha, x \rangle) - R'(\langle \alpha, y \rangle)| |\alpha|_V \leq Lip(R') |\langle \alpha, x - y \rangle| |\alpha|_V \\ \leq Lip(R') |\alpha|_V^2 |x - y|_{V'}.$$

so that Assumption 3.2-(3) holds true. Assumptions 2.2(ii) coupled with Remark 3.4 implies both Assumptions 3.2 (4) and that  $h_0^*$  is convex and Fréchet differentiable. The fact that  $h_0^*$  has Lipschitz differential is implied by (iii), so that also (5) holds true. Clearly (iv) implies (6). The last statement is straightforward.  $\square$

**Remark 3.6.** It is important to note that, in the case when the functions  $R$  and  $C$  are both quadratic, neither (1) nor (2) are satisfied in Assumption 3.3. Nonetheless necessary and sufficient conditions of optimality (see Theorem 4.5) hold true, and the value function results regular (see Remark 4.4 and Section 5.2.1 for details).  $\blacksquare$

#### 4. EQUILIBRIUM POINTS

Although the core of the section is the definition of equilibrium points of the abstract problem (P) and the investigation of their properties, some results are needed beforehand. Those obtained via Dynamic Programming, and contained in [44], are recalled for the reader's convenience in Section 4.1. On the other hand, Section 4.2 contains new material, and in particular a version of the Maximum Principle for problem (P). Finally Section 4.3 contains the analysis of equilibrium points.

**4.1. Dynamic Programming for problem (P).** We here recall the main results contained in [44]. If the value function is defined as

$$(4.1) \quad Z(t, x) = \inf_{u \in L_\lambda^p(t, +\infty; U)} J(t, x, u),$$

and, if one sets  $Z_0(x) := Z(0, x)$ , then  $Z(t, x) = e^{-\lambda t} Z_0(x)$ , so that the Hamilton–Jacobi–Bellman equation associated to the problem by means of Dynamic Programming reduces to that with initial time  $t = 0$ , that is

$$(4.2) \quad -\lambda\psi(x) + \langle \psi'(x), Ax \rangle - h_0^*(-B^*\psi'(x)) + g_0(x) = 0, \quad x \in H$$

(with  $\psi$  the unknown) whose candidate solution is  $Z_0(x)$ . We refer to  $p \mapsto h_0^*(-B^*p)$  as to the *Hamiltonian* function.<sup>7</sup>

**Definition 4.1.** *A function  $\psi$  is a classical solution of the stationary HJB equation (4.2) if it belongs to  $\Sigma_0(V')$  and satisfies (4.2) for every  $x \in D(A)$ .*

**Theorem 4.2.** *Let Assumptions 3.2 and 3.3 hold. Then there exists a unique classical solution  $\Psi$  to (4.2) and it is given by the value function of the optimal control problem, that is*

$$\Psi(x) = Z_0(x) = \inf_{u \in L_\lambda^p(0, +\infty; U)} J(0, x, u).$$

Once we have established that  $\Psi$  is the unique classical solution to the stationary HJB equation, and since  $\Psi$  is Fréchet differentiable with Lipschitz derivative, we can build optimal feedbacks and prove the following theorem.

**Theorem 4.3.** *Let Assumptions 3.2 and 3.3 hold. Let  $t \geq 0$  and  $x \in V'$  be fixed. Then there exists a unique optimal pair  $(u^*, y^*)$  at  $(t, x)$ . The optimal state  $y^*$  is the unique solution of the Closed Loop Equation*

$$(4.3) \quad \begin{cases} y'(\tau) = Ay(\tau) + B(h_0^*)'(-B^*\Psi'(y(\tau))), & \tau > t \\ y(t) = x \in V', \end{cases}$$

while the optimal control  $u^*$  is given by the feedback formula

$$u^*(s) = (h_0^*)'(-B^*\Psi'(y^*(s))).$$

where the optimal feedback map  $x \mapsto (h_0^*)'(-B^*\Psi'(x))$  is Lipschitz continuous.

**Remark 4.4.** There are relevant cases when Assumption 3.3 is not satisfied. One such example is the case, important for the applications, when costs  $g_0$  and  $h_0$  are quadratic (or linear + quadratic). Nonetheless Theorem 4.3 remains true, with identical proof to that provided in [44], if the value function  $Z_0$  is in  $C_{Lip}^1(V')$ . Indeed the regularity of  $Z_0$  implies that  $Z_0$  is a classical solution of the associated HJB equation (4.2) (to this extent see e.g. [63], ch. 6, Proposition 1.2, p. 225). In the case of quadratic costs, for instance, one proves that  $Z_0$  is itself quadratic, and hence in  $C_{Lip}^1(V')$ . Note also that if  $Z_0$  is not differentiable, then the closed loop equation (4.3) holds in the weaker sense of (4.8), as specified in the next section. ■

<sup>7</sup>Note that the function usually called Hamiltonian would be  $(p, x) \mapsto \langle p, Ax \rangle - h_0^*(-B^*p) + g_0(x)$ .

**4.2. Maximum Principle for Problem (P).** The results contained in this section, namely Theorems 4.5 and 4.6, are new to literature and add to the theory developed in [38, 40, 41, 43, 44]. They establish a Maximum Principle for the problem at hand, and connect it to the results on Dynamic Programming contained in those papers. The reader may find all of the proofs in the Appendix, as well as some additional results. We advise the reader that, differently from [44] and in view of Remark 4.4, the new results are proved avoiding Assumption 3.3. As a consequence, if on the one hand the regularity of the value function  $Z_0$  of (P) does not necessarily hold true, on the other hand we are able to treat the case of the limit exponent  $p = 2$ , and hence of quadratic costs  $g_0$  and  $h_0$ , so important for the applications.

In order to establish a maximum principle, we first need to define a dual system associated to the minimization problem. For all fixed  $x \in V'$  and  $t \geq 0$ , we consider the equation

$$(4.4) \quad \pi'(\tau) = (\lambda - A_1^*)\pi(\tau) - g'_0(y(\tau)), \quad \tau \in [t, +\infty)$$

where  $\pi : [t, +\infty) \rightarrow V$  (the dual variable, or co-state of the system) is the unknown, and  $y = y(\cdot; t, x, u)$  is the trajectory starting at  $x$  at time  $t$  and driven by control  $u$ , given by (3.1). We assume such equation is also subject to the following transversality condition

$$(4.5) \quad \lim_{T \rightarrow +\infty} e^{(\omega - \lambda)T} \pi(T) = 0.$$

When necessary, we denote any solution of (4.4)(4.5) also by  $\pi(\cdot; t, x, u)$  or by  $\pi(\cdot; t, x)$  to remark its dependence on the data.

Heuristically speaking, the candidate conditions of optimality associated to the problem are the following:

$$(4.6) \quad \begin{cases} y'(\tau) = Ay(\tau) + Bu(\tau), & \tau \geq t \\ y(t) = x \\ \pi'(\tau) = (\lambda - A_1^*)\pi(\tau) - g'_0(y(\tau)), & \tau \geq t \\ \lim_{T \rightarrow +\infty} e^{(\lambda - \omega)T} \pi(T) = 0, \\ -B^*\pi(\tau) \in \partial h_0(u(\tau)), & \tau \geq t. \end{cases}$$

The ODEs for  $y$  and  $\pi$  appearing in (4.6) are intended, as it is usual in these cases, in *mild* sense, see Definition A.1 in Appendix A. Moreover, by conjugation formula, we have

$$(4.7) \quad -B^*\pi(\tau) \in \partial h_0(u(\tau)) \iff u(\tau) = (h_0^*)'(-B^*\pi(\tau)).$$

We refer to (4.7) as to *maximum condition*. It has to be satisfied for a.a.  $\tau \geq t$ .

The conditions listed in (4.6) prove to be necessary and sufficient for optimality for all  $p \geq 2$ , in the sense specified next.

**Theorem 4.5. (Maximum Principle).** *Let Assumptions 3.2 be satisfied. Let  $p \geq 2$ ,  $q = \frac{p}{p-1}$  and  $\lambda > (2\omega) \vee \omega$ . Let  $t \geq 0$ ,  $x \in V'$ .*

- (i) Let  $(u, y) \in L_\lambda^p(t, +\infty; U) \times L_{loc}^1(t, +\infty; V')$  be a given admissible pair at  $(t, x)$ . If there exists a function  $\pi \in L_\lambda^q(t, +\infty; V)$  satisfying, along with  $u$  and  $y$ , the system (4.6), then  $(u, y)$  is optimal at  $(t, x)$  for the problem of minimizing (3.1)(3.2).
- (ii) Assume further that, either  $p > 2$  and  $\lambda > 0$ , or  $p = 2$ ,  $\lambda > 0$  and  $\omega < 0$ . Then the viceversa of (i) holds, i.e., any couple  $(u^*, y^*)$  optimal at  $(t, x)$  necessarily admits a costate  $\pi \in L_\lambda^q(t, +\infty; V)$  satisfying, along with  $u^*$  and  $y^*$ , system (4.6).

The next theorem contains the so-called *co-state inclusion*. Note that the case of  $p = 2$  is discussed separately, as the value function is not necessarily Fréchet differentiable (unless Assumption 3.3 holds or ad hoc regularity results are given).

**Theorem 4.6. (Co-state inclusion).** *In Assumptions 3.2, for  $\lambda > \max\{0, \omega, 2\omega\}$ , suppose that either  $p > 2$ , or  $p = 2$  and  $\omega < 0$ . Let  $(u^*, y^*)$  be optimal at  $(t, x) \in [0, +\infty) \times V'$ , and let  $\pi^*(\cdot; t, x) \in L_\lambda^q(t, +\infty; V)$  be the associated co-state. Let also  $Z_0$  be the value function of problem (P). Then*

$$\pi^*(\tau; t, x) = \pi^*(\tau; \tau, y^*(\tau)) \in \partial Z_0(y^*(\tau)), \quad \forall \tau \geq t.$$

where  $\partial Z_0$  is the subdifferential of the convex function  $Z_0$ . If in addition  $p > 2$ , then  $Z_0 \in \Sigma_0(V')$  and  $Z_0$  coincides with  $\Psi$ , so that

$$\pi^*(\tau; t, x) = \pi^*(\tau; \tau, y^*(\tau)) = \Psi'(y^*(\tau)), \quad \forall \tau \geq t.$$

**Remark 4.7.** Note that, for  $p \geq 2$  and  $\lambda > 0$ , and by making use of Theorem 4.5, and of equations (4.6) and (4.7), one obtains

$$y'(\tau) = Ay(\tau) + B(h_0^*)'(-B^*\pi(\tau)), \quad \tau \geq t$$

so that the general version of the closed loop equation (4.3) becomes a differential inclusion

$$(4.8) \quad y'(\tau) \in Ay(\tau) + B(h_0^*)'(-B^*\partial Z_0(y(\tau))), \quad \tau \geq t.$$

also to be intended in mild sense. ■

**4.3. Equilibrium points.** We give two different definitions of equilibrium points for problems (P), and later show to which extent they are equivalent.

**Definition 4.8.** *A MP-equilibrium point of problem (P) is any stationary solution  $(x, \pi, u) \in V' \times V \times U$  of (4.6). This is equivalent to require that  $(x, \pi, u)$  belongs to  $D(A) \times D(A_1^*) \times U$  and satisfies*

$$(4.9) \quad \begin{cases} Ax + Bu = 0 \\ (\lambda - A_1^*)\pi - g'_0(x) = 0, \\ u = (h_0^*)'(-B^*\pi). \end{cases}$$

*A CLE-equilibrium point of problem (P) is any  $x \in V'$  that is a stationary solution of the closed loop equation (4.8). This is equivalent to require  $x \in D(A)$  and*

$$(4.10) \quad Ax + B(h_0^*)'(-B^*\partial Z_0(x)) \ni 0.$$

**Remark 4.9.** When  $0 \in \rho(A)$  and  $\lambda \in \rho(A_1^*)$ , then (4.9) is equivalent to

$$(4.11) \quad \begin{cases} x = -A^{-1}Bu \\ \pi = (\lambda - A_1^*)^{-1}g'_0(x), \\ u = (h_0^*)'(-B^*\pi). \end{cases}$$

and (4.10) is equivalent to

$$(4.12) \quad x \in -A^{-1}B(h_0^*)'(-B^*\partial Z_0(x)).$$

As a consequence of Remark 4.4, the equations (4.8), (4.10) and (4.12) hold as equalities with  $\Psi'(x)$  in place of  $\partial Z_0(x)$  when  $Z_0$  Fréchet differentiable in  $V'$  (e.g. when  $p > 2$ , or when regularity can be proven separately). ■

The proof of the equivalences in the above definition is straightforward as they are based on standard regularity of convolutions of semigroups. We omit them for brevity.

We have the following result.

**Theorem 4.10.** *Let Assumptions 3.2 be satisfied,  $p \geq 2$ ,  $\lambda > (2\omega) \vee \omega$ .*

- (i) *Let  $(\bar{x}, \bar{\pi}, \bar{u}) \in D(A) \times D(A_1^*) \times U$  be any MP-equilibrium point. Then the constant control  $\bar{u}$  is optimal at  $(0, \bar{x})$  and*

$$(4.13) \quad A\bar{x} + B(h_0^*)'(-B^*(\lambda - A_1^*)^{-1}g'_0(\bar{x})) = 0,$$

*moreover  $\bar{x}$  is a CLE-equilibrium point and*

$$(4.14) \quad \partial Z_0(\bar{x}) \ni (\lambda - A_1^*)^{-1}g'_0(\bar{x}).$$

- (ii) *Let  $\hat{x} \in D(A)$  be a CLE-equilibrium point,  $\lambda > 0$ . Let either  $p > 2$ , or  $p = 2$  and  $\omega < 0$ . Assume that  $Z_0$  is Fréchet differentiable in  $V'$ . Then  $(\hat{x}, \hat{\pi}, \hat{u})$ , where*

$$\hat{\pi} := (\lambda - A_1^*)^{-1}g'_0(\hat{x}) \quad \text{and} \quad \hat{u} := (h_0^*)'(-B^*\hat{\pi}),$$

*is an MP-equilibrium point, the control  $\hat{u}$  is optimal at  $(0, \hat{x})$  and  $Z'_0(\hat{x}) = \hat{\pi} = (\lambda - A_1^*)^{-1}g'_0(\hat{x})$ .*

One important consequence of the above theorem is that it provides the following equation for a CLE-equilibrium point (or for the first component of an MP-equilibrium point)

$$(4.15) \quad Ax + B(h_0^*)'(-B^*(\lambda - A_1^*)^{-1}g'_0(x)) = 0.$$

In addition, whenever  $0 \in \rho(A)$  (this assumption is satisfied in the optimal investment problem with vintage capital described in Section 2) solutions of (4.15) can be regarded as fixed points of the operator  $T : V' \rightarrow V'$ , defined by

$$(4.16) \quad Tx := -A^{-1}B(h_0^*)'(-B^*(\lambda - A_1^*)^{-1}g'_0(x)).$$

For the applications, the most efficient way of making use of such relations is to rewrite them in terms of the specific sets of data, and compute when possible the optimal equilibrium distributions. In particular, in Section 5 we will see how (4.16) is interpreted in

terms of the data of optimal investment with vintage capital, so that fixed points of  $T$  may be directly computed by solving a numeric equation.

However, in the general case, it is possible to provide sufficient conditions for the existence and uniqueness of a fixed point of the operator  $T$  using well known fixed point theorems although, as one expects, such conditions may hardly be very sharp. To this extent, we provide here only Lemma 4.11, which is a straightforward application of the contraction mapping principle.

**Lemma 4.11.** *Let Assumptions of Theorem 4.10 be satisfied. Assume moreover that*

$$\lambda - \omega > \|(A)^{-1}\|_{\mathcal{L}(V')} \|B\|_{\mathcal{L}(U,V')}^2 [(h_0^*)'] [g_0'].$$

*Then there exists a unique solution  $\bar{x} \in D(A)$  to the equation (4.16).*

**Remark 4.12.** The operator  $T$  above is considered as an operator from  $V'$  to itself. Since its image is contained in  $D(A)$ , when looking for fixed points, it is also equivalent to look at it as an operator from  $D(A)$  to itself, considered as a subspace of  $V'$ , as done in Lemma 5.4. ■

**Remark 4.13.** All above results could be generalized to the case in which we have state constraints and the function  $g_0$  is convex but not necessarily Fréchet differentiable. This could be done using the results of [42] and generalizing them to the infinite horizon case, using the same arguments in [53]. Clearly, at points where  $g_0$  is not Fréchet differentiable, one would have to choose an element of the subdifferential of  $g_0$ . ■

**4.4. Stability.** Once existence (and possibly uniqueness) of equilibrium points is proven, it is possible to study their stability properties adapting known results such as those in [59] or in chapter 9 in [64], or by direct proof, as we see next. In all cases, stability will be proven with respect to the topology of  $V'$ . For the reader's convenience we recall the definition here below.

**Definition 4.14.** *A CLE-equilibrium point  $\bar{x} \in D(A)$  is stable in the topology of  $V'$  if,  $\forall \epsilon > 0, \exists \delta > 0$  such that, if  $x \in V'$  and  $x^*(\cdot)$  is the optimal trajectory starting at  $x$ , then  $|x - \bar{x}|_{V'} < \delta \Rightarrow |x^*(t) - \bar{x}|_{V'} < \epsilon$ . If in addition  $\lim_{t \rightarrow \infty} |x^*(t) - \bar{x}|_{V'} = 0$  then  $\bar{x}$  is asymptotically stable. Finally, if the same property hold true for all  $x \in V'$ , then  $\bar{x}$  is globally asymptotically stable.*

The first criterium to establish stability is contained in the following proposition and makes use of the linearization method. The proof follows from Corollary 2.2 in [59].

**Proposition 4.15.** (Stability by linearization) *Let Assumption 3.2 be satisfied and  $\Psi \in C_{Lip}^1$ . For  $x \in V'$  set  $f(x) := B(h_0^*)'(-B^*\Psi'(x))$ , and assume that  $\bar{x} \in D(A)$  is a CLE-equilibrium point for (P), that  $f$  is continuously Fréchet differentiable at a neighborhood of  $\bar{x}$ , and denote by  $\sigma_{\bar{x}}$  the spectrum of the operator  $A + f'(\bar{x})$ . If  $\sup(\text{Re}\sigma_{\bar{x}}) < 0$ , then  $\bar{x}$  is stable in the topology of  $V'$ . Moreover, it is also asymptotically stable in the topology of  $V'$ . If  $\sup(\text{Re}\sigma_{\bar{x}}) > 0$ ,  $\bar{x}$  is unstable in the topology of  $V'$ .*

Another result that can be used is the following. Recall that  $\langle \cdot, \cdot \rangle_{V'}$  indicates the inner product in  $V'$ .

**Proposition 4.16.** (Stability by dissipativity) *Let Assumption 3.2 be satisfied and let  $\Psi \in C_{Lip}^1(V')$ . For  $x \in V'$ , and  $f(x) = B(h_0^*)'(-B^*\Psi'_0(x))$ , and assume that, when  $x \in D(A)$ , the solution of the closed loop equation*

$$(4.17) \quad \begin{cases} y'(t) = Ay(t) + f(y(t)), & t > 0 \\ y(0) = x, \end{cases}$$

*belongs to  $D(A)$  for all  $t \geq 0$ . Let  $\bar{x} \in D(A)$  be a CLE-equilibrium point for (P). Assume that  $A + f$  is dissipative near  $\bar{x}$ , i.e. there exists an open ball  $I(\bar{x})$  in  $V'$  centered at  $\bar{x}$ , and  $\xi \leq 0$  such that, for every  $x \in I(\bar{x}) \cap D(A)$ ,*

$$(4.18) \quad (A(x - \bar{x}) + f(x) - f(\bar{x}))|x - \bar{x}\rangle_{V'} \leq \xi|x - \bar{x}|_{V'}^2.$$

*Then  $\bar{x}$  is stable in the topology of  $V'$ . If  $\xi < 0$  then  $\bar{x}$  is asymptotically stable in the topology of  $V'$ . If  $A + f$  is dissipative on the whole  $V'$  and  $\xi < 0$  then  $\bar{x}$  is globally asymptotically stable.*

**Corollary 4.17.** *Let the assumptions of Proposition 4.16 be verified, except (4.18). Let the operator  $A$  satisfy  $(Ax|x)_{V'} \leq -\theta|x|_{V'}^2$ , for all  $x$  in  $V'$ , with  $\theta > 0$  a fixed constant. If there exists a neighborhood  $I$  of  $\bar{x}$  where  $f$  is Lipschitz continuous (in the topology of  $V'$ ) with Lipschitz constant strictly smaller than  $\theta$ , then  $\bar{x}$  is asymptotically stable in the topology of  $V'$ . If  $f$  is Lipschitz continuous in  $V'$  with Lipschitz constant strictly smaller than  $\theta$  then  $\bar{x}$  is globally asymptotically stable.*

In particular, the above corollary may be applied to the examples in Section 5, see e.g. subsection 5.2.1.

## 5. APPLICATION TO OPTIMAL INVESTMENT WITH VINTAGE CAPITAL

The aim of this section is to show how valuable our general theory can be when analyzing specific applications, and in particular when trying to derive analytic formulas for equilibrium distributions. This process unfolds by computing MP-equilibrium/CLE-equilibrium points for that problem rephrased in abstract form as in Section 3.3.

We begin by noting that the value function  $V(t, x) = e^{-\lambda t}V(0, x) = e^{-\lambda t}V_0(x)$  of the optimal control problem described in Section 2 (see (2.5)), satisfies, as a consequence of (2.4), (3.9) and (4.1),

$$V_0(x) = -Z_0(x)$$

where  $Z_0$  is the value function, defined in Section 4.1, of the abstract problem (P) for  $t = 0$ . Note that  $V_0$  is a concave function, as  $Z_0$  is convex. Note also that, under additional assumptions (e.g. Assumption 3.3, or regularity assumptions on  $Z_0$ ),  $Z_0$  is

the unique classical solution of HJB equation (4.2) in the sense of Definition 4.1. As a consequence, the natural co-state for the maximization problem would be

$$\zeta(\tau, s) = -\pi(\tau)[s],$$

with  $\pi$  the co-state of the abstract problem whose properties are described in Theorem 4.5 and 4.6. Then the optimality conditions (4.6) for a triplet  $(K^*, \zeta^*, u^*)$  can be written as the following set of equations

$$(5.1) \quad u_0^*(\tau) = (C_0^*)'(\zeta^*(\tau, 0)), \quad u_1^*(\tau, s) = ((C_1^*)'(\zeta^*(\tau, \cdot)))[s] = [c_1(s, \cdot)]'(\zeta^*(\tau, s))$$

$$(5.2) \quad \zeta^*(\tau, s) = \int_s^{\bar{s}} e^{-(\lambda+\mu)(\xi-s)} R' \left( \int_0^{\bar{s}} \alpha(\theta) K^*(\tau + \xi - s, \theta) d\theta \right) \alpha(\xi) d\xi;$$

$$(5.3) \quad \lim_{T \rightarrow +\infty} e^{(\lambda-\omega)T} \zeta(T, s) = 0, \quad a.a. \ s \in [0, \bar{s}].$$

(5.4)

$$K^*(\tau, s) = \begin{cases} e^{-\mu(\tau-s)} x(s - \tau + t) + \int_0^{\tau-t} e^{-\mu\sigma} u_1^*(\tau - \sigma, s - \sigma) d\sigma & s \in [\tau - t, \bar{s}], \ \tau \in [t, \bar{s} + t] \\ e^{-\mu(\tau-s)} u_0^*(\tau - s) + \int_0^s e^{-\mu\sigma} u_1^*(\tau - \sigma, s - \sigma) d\sigma & s \in [0, \tau - t], \ \tau \in [t, \bar{s} + t] \\ 0 & s \in [0, \bar{s}], \ \tau \in (\bar{s} + t, +\infty) \end{cases}$$

Note that (5.1) and (5.2) are derived from (4.6) by making use of (4.7) and (A.15), while (5.4) is well known and can be obtained by means of characteristics method (see e.g. [14]). The following proposition is an immediate consequence of Theorem 4.5 and of Lemma 3.5.

**Proposition 5.1.** *Under Assumptions 2.2, with  $p \geq 2$  and  $\lambda > 0$ , the optimality conditions (5.1)(5.2)(5.4) are necessary and sufficient for a couple  $(u^*, K^*)$  to be optimal at  $x$  for the problem of optimal investment with vintage capital described in Section 2.*

**5.1. Characterization of Equilibrium Points.** It is natural to define an equilibrium point for the problem consistently with Section 4.3. For the reader's convenience, Definition 4.8 is reformulated below in terms of the problem of optimal investment with vintage capital.

It is important to note that an equilibrium point is actually a function of the variable  $s$  (although independent of  $t$ ) dependent on the variable  $s$ , hence an *equilibrium distribution*.

**Definition 5.2.** *In reference to the the optimal investment problem with vintage capital:*

- (i) a MP-equilibrium point is a stationary solution  $(x, \zeta, (u_0, u_1)) \in L^2(0, \bar{s}) \times H^2(0, \bar{s}) \times (\mathbb{R} \times L^2(0, \bar{s}))$  of the system of equations (5.1)(5.2)(5.4);
- (ii) for  $V_0$  Fréchet differentiable, a CLE-equilibrium point is any  $x \in L^2(0, \bar{s})$  which is a stationary solution of equation (5.4) when

$$u_0^*(\tau) = [(C_0^*)'(V_0'(K^*(\tau, \cdot)))] [0], \quad u_1^*(\tau, s) = [(C_1^*)'(V_0'(K^*(\tau, \cdot)))] [s].$$

**Remark 5.3.** Several remarks are here due.

- (1) Theorem 4.10 implies, in the assumptions of Proposition 5.1 that (i) and (ii) are equivalent in the following sense: the first component  $x$  of a MP-equilibrium point  $(x, p, (u_0, u_1))$  is also a CLE-equilibrium point; conversely, when  $V_0$  is Fréchet differentiable, a CLE-equilibrium point  $x$  can be used to build a MP-equilibrium point by means of (5.1)(5.2)(5.4), having  $x$  as first component.
- (2) If  $V_0$  is not Fréchet differentiable, the closed loop equation (as well as the definition above) may be generalized to a differential inclusion in the sense of (4.8), where  $V'_0$  is replaced by the superdifferential  $\partial V_0$ .
- (3) Definition 5.2 is consistent with Definition 4.8 as here  $D(A) = L^2(0, \bar{s})$  and  $D(A_1^*) \subseteq H^2(0, \bar{s})$ .

■

We further characterize MP-equilibrium/CLE-equilibrium points as fixed points of a suitable operator. To this extent we define

$$(5.5) \quad \bar{\alpha}(s) = \int_s^{\bar{s}} e^{-(\mu+\lambda)(\sigma-s)} \alpha(\sigma) d\sigma$$

*i.e.*  $\bar{\alpha}(s)$  is the *discounted return* associated with a unit of capital of vintage  $s$ .

**Lemma 5.4.** *Under Assumptions 2.2,  $\bar{x}$  is a CLE-equilibrium point if and only if it is a fixed point of the operator  $T : L^2(0, \bar{s}) \rightarrow L^2(0, \bar{s})$  defined by*

$$(5.6) \quad (Tx)[s] = (C_0^*)' (R'(\langle \alpha, x \rangle) \bar{\alpha}(0)) e^{-\mu s} + \int_0^s e^{-\mu(s-\sigma)} [c_1(\sigma, \cdot)^*]' (R'(\langle \alpha, x \rangle) \bar{\alpha}(\sigma)) d\sigma,$$

*that is, if and only if  $(T\bar{x})[s] = \bar{x}(s)$  for a.e.  $s$  in  $[0, \bar{s}]$ . Moreover  $(\bar{x}, \bar{\zeta}, (\bar{u}_0, \bar{u}_1))$  where*

$$(5.7) \quad \bar{\zeta}(s) = R'(\langle \alpha, \bar{x} \rangle) \bar{\alpha}(s), \quad \bar{u}(s) = (h_0^*)' (B^* \bar{\zeta})(s), \quad \text{for a.e. } s \in [0, \bar{s}]$$

*is a MP-equilibrium point.*

The proof of the lemma is contained in Appendix C.

Note that solving the equation  $Tx = x$  within a space of functions is not particularly handy. Nonetheless solving such functional equation is equivalent - and in the generality of cases - to solving a *numeric equation*. In this sense, the following theorem contains the most interesting result of the section.

**Theorem 5.5.** *Let Assumptions 2.2 be satisfied, and let  $T$  be given by (5.6). Moreover, for any  $\eta \in \mathbb{R}$  and  $s \in [0, \bar{s}]$ , consider the function*

$$(5.8) \quad F(\eta)[s] = (C_0^*)' (\eta \bar{\alpha}(0)) e^{-\mu s} + \int_0^s e^{-\mu(s-\sigma)} [c_1(\sigma, \cdot)^*]' (\eta \bar{\alpha}(\sigma)) d\sigma.$$

*Then  $\bar{x} \in L^2(0, \bar{s})$  is a solution of  $Tx = x$ , if and only if*

$$(5.9) \quad \bar{x}(s) = F(\bar{\eta})[s]$$

*with  $\bar{\eta}$  a solution in  $\mathbb{R}$  of*

$$(5.10) \quad \eta = R'(\langle \alpha, F(\eta) \rangle).$$

The proof of the theorem is contained in Appendix C.

**Remark 5.6.** Note that the solution of (5.10) is unique and nonnegative when, for instance,  $R'(\langle \alpha, F(0) \rangle) \geq 0$ . Indeed  $R'$  is decreasing and  $C_0^*$  and  $c_1(\sigma, \cdot)^*$  are convex functions, then the right hand side of (5.10) is a positive decreasing function of  $\eta$ . Hence the function

$$(5.11) \quad \theta(\eta) := \eta - R'(\langle \alpha, F(\eta) \rangle)$$

satisfies  $\theta(0) \leq 0$ , is strictly increasing to  $+\infty$ , and hence has exactly one nonnegative zero.  $\blacksquare$

Note that (5.6) and (5.9) are general formulas, holding for any choice of costs and revenues, as long as they satisfy Assumptions 2.2. More explicit formulas for the optimal equilibrium distribution may be derived once costs and revenues are further specified.

**5.2. Linear-quadratic costs.** Now we make formulas more explicit in the case of cost functions satisfying (2.7). We derive

$$(5.12) \quad C_0^*(p_0) = \frac{(p_0 - q_0)^2}{4\beta_0}, \quad C_1^*(p_1) = \int_0^{\bar{s}} \frac{(p_1(s) - q_1(s))^2}{4\beta_1(s)} ds$$

so that

$$(C_0^*)'(p_0) = \frac{p_0 - q_0}{2\beta_0}, \quad (C_1^*)'(p_1)(s) = \frac{p_1(s) - q_1(s)}{2\beta_1(s)}$$

In this case, (5.6) becomes

$$(5.13) \quad \begin{aligned} (Tx)[s] &= \frac{R'(\langle \alpha, x \rangle) \bar{\alpha}(0) - q_0}{2\beta_0} e^{-\mu s} + \int_0^s e^{-\mu(s-\sigma)} \frac{R'(\langle \alpha, x \rangle) \bar{\alpha}(\sigma) - q_1(\sigma)}{2\beta_1(\sigma)} d\sigma \\ &= R'(\langle \alpha, x \rangle) w_1(s) - w_2(s) \end{aligned}$$

where  $w_1$  and  $w_2$  are the positive functions

$$(5.14) \quad w_1(s) = \frac{\bar{\alpha}(0)}{2\beta_0} e^{-\mu s} + \int_0^s e^{-\mu(s-\sigma)} \frac{\bar{\alpha}(\sigma)}{2\beta_1(\sigma)} d\sigma$$

$$(5.15) \quad w_2(s) = \frac{q_0}{2\beta_0} e^{-\mu s} + \int_0^s e^{-\mu(s-\sigma)} \frac{q_1(\sigma)}{2\beta_1(\sigma)} d\sigma.$$

If in addition we define the positive coefficients

$$(5.16) \quad c_1 = \langle w_1, \alpha \rangle = \int_0^{\bar{s}} \alpha(s) w_1(s) ds, \quad c_2 = \langle w_2, \alpha \rangle = \int_0^{\bar{s}} \alpha(s) w_2(s) ds.$$

then the following result follows as a consequence of Theorem 5.5.

**Corollary 5.7.** *Let Assumption 2.2 and (2.7) be satisfied. Let  $w_1$ ,  $w_2$ ,  $c_1$ , and  $c_2$  be defined respectively by (5.14), (5.15) and (5.16). Then  $(\bar{x}, \bar{\zeta}, \bar{u})$  is a MP-equilibrium point if and only if  $\eta \in \mathbb{R}$  is a solution of*

$$(5.17) \quad \eta = R'(\eta c_1 - c_2).$$

and

$$\bar{x}(s) = -w_2(s) + \eta w_1(s).$$

and moreover  $\bar{\zeta}$  and  $\bar{u}$  are given by (5.7). The constant control  $\bar{u}$  is optimal at  $\bar{x}$ ,  $\bar{x}$  is also a CLE-equilibrium point and, if  $V_0$  is Fréchet differentiable, then it is also the unique CLE-equilibrium point.

**Remark 5.8.** If the CLE-equilibrium point  $\bar{x}$  identified by Corollary 5.7 is such that  $\bar{x}(s) \geq 0$  at all  $s$ , then  $\bar{x}$  is also a CLE-equilibrium point for the problem with state constraints  $k(\tau, s) \geq 0$  for all  $s$  and  $\tau$  (see also Remarks 2.1 and 3.1). ■

**Remark 5.9.** Note that Assumptions 2.2 are satisfied here with  $p = 2$ , so that  $V_0$  is not necessarily Fréchet differentiable. That implies that, although the first component  $\bar{x}$  of a MP-equilibrium point  $(\bar{x}, \bar{\zeta}, \bar{u})$  is also a CLE-equilibrium point, the viceversa may fail: there may be CLE-equilibrium points which do not derive as first components of a MP-equilibrium point, i.e. solutions of the stationary closed loop equation which fail to be optimal. For a further discussion on regularity of  $V_0$ , the reader is referred to Section 5.2.1. ■

Once  $R$  is chosen, the results in Corollary 5.7 leads to an explicit formula for that CLE-equilibrium point, as illustrated in the next lemma.

**Lemma 5.10.** *In the assumptions of Corollary 5.7, there exists a unique CLE-equilibrium point  $\bar{x}$ , described by the formuals below, for the associated choices of the revenue  $R$ :*

(i) *If  $R(Q) = -aQ^2 + bQ$ , then*

$$\bar{x} = -w_2 - \frac{2ac_2 + b}{1 + 2ac_1} w_1;$$

(ii) *If  $R(Q) = \ln(1 + Q)$ , for  $Q \geq 0$  and  $R(Q) = Q$  for  $Q < 0$ , then*

$$\bar{x} = -w_2 + \frac{\sqrt{(1 - c_2)^2 + 4c_1} - (1 - c_2)}{2c_1} w_1$$

(iii) *If  $R(Q) = b[(\nu + Q)^\gamma - \nu]$ , with  $\gamma \in (0, 1)$ ,  $b, \nu > 0$ , for  $Q \geq 0$  and  $R(Q) = \gamma\nu^{\gamma-1}Q$  for  $Q < 0$ , then  $\bar{x} = -w_2 + \bar{\eta}w_1$  where  $\bar{\eta}$  is the unique positive solution of*

$$\eta = \frac{b\gamma}{(\nu + c_1\eta - c_2)^{1-\gamma}}.$$

(iv) *If  $R(Q) = bQ^\gamma$ , with  $\gamma \in (0, 1)$ ,  $b > 0$ , for  $Q \geq 0$ , and  $R(Q) = -\infty$  for  $Q < 0$  (case with state constraints) then  $\bar{x} = -w_2 + \bar{\eta}w_1$  where  $\bar{\eta}$  is the unique positive solution of*

$$\eta = \frac{b\gamma}{(c_1\eta - c_2)^{1-\gamma}}.$$

*Proof.* The proof follows from straightforward computations. □

5.2.1. *Stability of Equilibrium Distributions.* We close the section on linear-quadratic costs (2.7) by briefly discussing stability of equilibrium distributions and, in some sub-cases, the regularity of the value function, by applying the results contained in Section 4.4. The concept of stability here used is that of Definition 4.14, which is natural in this context. We remark though that the convergence of functions there mentioned (i.e. in the topology of  $V'$ ) is *not* a convergence in the space  $L^2(0, \bar{s})$  but, roughly speaking, the (weaker) convergence of their primitive functions.

**Lemma 5.11.** *In the assumptions of Corollary 5.7, suppose in addition that the value function  $V_0$  is Fréchet differentiable, and set  $\xi = -\mu + [V_0]_L |\delta_0|_{V'}^2 / (4\beta_0)$ , where  $[V_0]_L$  indicates the Lipschitz constant of the gradient  $V'_0$ . If  $\xi \leq 0$  (respectively,  $\xi < 0$ ) then  $\bar{x}$  is stable (resp., asymptotically stable) in the sense of Definition 4.14.*

The proof of the lemma is contained in Appendix C.

**Remark 5.12.** In particular, the previous Lemma applies when  $R$  is of the type described in Lemma 5.10 (i). Indeed with some extra work one shows that in this case the value function of the abstract problem is of type

$$\Psi(x) = \langle Cx, x \rangle + \langle d, x \rangle + e,$$

for a suitable linear operator  $C : V' \rightarrow V$ ,  $d \in V$  and  $e \in \mathbb{R}$ , where  $V$  and  $V'$  are the spaces introduced in Subsection 3.3. Hence  $\Psi$  is differentiable with Fréchet differential  $\Psi'(x) = Cx + d$ , and  $[\Psi]_L = \|C\|_{\mathcal{L}(V',V)}$  (for a proof, we refer the reader to [60], vol 1, ch.2). This applies in particular to the linear-quadratic examples of Section 6. ■

5.3. **Linear-quadratic costs, constrained control.** We now choose costs as in (2.8). We then derive

$$g_{\beta,M}^*(v) = \sup_{|u| \leq M} \{vu - \beta u^2\} = \begin{cases} \frac{v^2}{4\beta} & |v| \leq 2\beta M \\ M|v| - \beta M^2 & \text{otherwise} \end{cases}$$

Note that  $g_{\beta,M}^*$  is a  $C^1$  function, with Lipschitz derivative

$$(g_{\beta,M}^*)'(v) = \begin{cases} \frac{v}{2\beta} & |v| \leq 2\beta M \\ M & v > 2\beta M \\ -M & v < -2\beta M \end{cases}$$

As a consequence, the Legendre transform of  $C$  is

$$C^*(v) = g_{\beta_0, M_0}^*(\beta_0 - q_0) + \int_0^{\bar{s}} g_{\beta_1(s), M_1}^*(v_1(s) - q_1(s)) ds$$

which is Fréchet differentiable with differential

$$(C^*)'(v)(s) = \left( (g_{\beta_0, M_0}^*)'(v_0 - q_0); (g_{\beta_1(s), M_1}^*)'(v_1(s) - q_1(s)) \right)$$

while the operator (5.6) is given by

$$T^M x(s) = (g_{\beta_0, M_0}^*)' (R'(\langle \alpha, x \rangle) \bar{\alpha}(0) - q_0) e^{-\mu s} + \int_0^s e^{-\mu(s-\sigma)} (g_{\beta_1(\sigma), M_1}^*)' (R'(\langle \alpha, x \rangle) \bar{\alpha}(\sigma) - q_1(\sigma)) d\sigma$$

**Remark 5.13.** Note that, with this choice of costs  $C$ , Assumptions 2.2 are satisfied with  $p > 2$  so that, by Theorem 4.2, the value function  $V_0$  is in  $C_{Lip}^1$ . By Lemma 5.4 and Theorem 5.5 we then get that there exists a unique CLE-equilibrium point, coinciding with the first component of the unique MP-equilibrium point. ■

From this point on, one may proceed as in the proof of Theorem 5.5 and Lemma 5.10 and compute CLE-equilibrium points, once the data  $\alpha, q_1, q_0$  are further specified.

**5.4. Power costs.** We now choose costs as in (2.10) and set  $q = \frac{p}{p-1}$ . Note that  $p > 2$  implies  $q \in (1, 2)$ . The convex conjugate of the costs are then

$$(5.18) \quad f_{\beta}^*(v) = \begin{cases} (\beta p)^{1-q} q^{-1} v^q - \theta v + \beta \theta^p & v \geq \beta p \theta^{p-1} \\ 0 & v < \beta p \theta^{p-1} \end{cases}$$

with Lipschitz derivative

$$(5.19) \quad (f_{\beta}^*)'(v) = \begin{cases} (\beta p)^{1-q} v^{q-1} - \theta & v \geq \beta p \theta^{p-1} \\ 0 & v < \beta p \theta^{p-1}. \end{cases}$$

As a consequence the Legendre transform of  $C$  is

$$C^*(v) = f_{\beta_0}^*(v_0 - q_0) + \int_0^{\bar{s}} f_{\beta_1(s)}^*(v_1(s) - q_1(s)) ds,$$

which is a  $C^1$  function with Lipschitz differential

$$(C^*)'(v)(s) = \left( (f_{\beta_0}^*)'(v_0 - q_0); (f_{\beta_1(s)}^*)'(v_1(s) - q_1(s)) \right).$$

Moreover

$$T^{\theta} x(s) = (f_{\beta_0}^*)' (R'(\langle \alpha, x \rangle) \bar{\alpha}(0) - q_0) e^{-\mu s} + \int_0^s e^{-\mu(s-\sigma)} (f_{\beta_1(\sigma)}^*)' (R'(\langle \alpha, x \rangle) \bar{\alpha}(\sigma) - q_1(\sigma)) d\sigma.$$

**Remark 5.14.** Note that Remark 5.13 applies also to this case. ■

## 6. SENSITIVITY ANALYSIS IN TWO SPECIAL CASES

We here analyze further the case of linear-quadratic costs discussed in Section 5.2, and develop sensitivity analysis accordingly. In particular we assume

$$(6.1) \quad \alpha(s) \equiv \alpha, \quad \beta_1(s) \equiv \beta_0, \quad q_1(s) = q_0 e^{-ws}.$$

Summing up, the objective functional of the profit maximizing firm is

$$(6.2) \quad \int_0^\infty e^{-\lambda t} \left( R(Q(K(t))) - \int_0^{\bar{s}} \left( q_1(s) u_1(t, s) + \frac{1}{2} \beta_0 u_1^2(t, s) \right) ds \right) dt \\ - \int_0^\infty e^{-\lambda t} \left( q_0 u_0(t) + \frac{1}{2} \beta_0 u_0^2(t) \right) dt.$$

We study separately the cases of linear-quadratic and power revenues, depicted respectively in Lemma 5.10 (i) and (iii).

**6.1. Linear-Quadratic Revenues.** We here assume

$$(6.3) \quad R(Q) = bQ - aQ^2$$

as in Lemma 5.10 (i), so that the equilibrium distribution there described equals

$$(6.4) \quad K^*(s) = -w_2(s) + \eta w_1(s),$$

in which

$$(6.5) \quad w_1(s) = \frac{\alpha}{2\beta_0(\mu + \lambda)} \left( e^{-\mu s} - e^{-\mu s} e^{-(\mu+\lambda)\bar{s}} + \frac{1 - e^{-\mu s}}{\mu} + \right. \\ \left. - \frac{e^{-(\mu+\lambda)\bar{s}}}{2\mu + \lambda} (e^{(\mu+\lambda)s} - e^{-\mu s}) \right),$$

$$(6.6) \quad w_2(s) = \frac{q_0}{2\beta_0(\mu - w)} (e^{-ws} - e^{-\mu s} (1 - \mu + w)).$$

and

$$\eta = \frac{b - 2ac_2}{1 + 2ac_1}$$

where

$$(6.7) \quad c_1 = \frac{\alpha^2}{2\beta_0(\mu + \lambda)} \left( \frac{1 - \mu}{\mu^2} (e^{-\mu\bar{s}} - 1) + \frac{2\mu + \lambda - 1}{\mu(2\mu + \lambda)} e^{-(2\mu+\lambda)\bar{s}} \right. \\ \left. + \frac{\bar{s}}{\mu} - \frac{1}{(2\mu + \lambda)(\mu + \lambda)} + \frac{1 - (\mu + \lambda)}{\mu(\mu + \lambda)} e^{-(\mu+\lambda)\bar{s}} \right),$$

$$(6.8) \quad c_2 = \frac{\alpha q_0}{2\beta_0(\mu - w)} \left( \frac{\mu - w - 1}{\mu} (1 - e^{-\mu\bar{s}}) + \frac{1}{w} (1 - e^{-w\bar{s}}) \right).$$

The explicit expression for the equilibrium distribution capital stock for every age,  $K^*(s)$ , allows us to obtain interesting economic implications. To illustrate, we establish some numerical results, which mostly are analytically proved as well. We start out from the following parameter values:

$$(6.9) \quad \alpha = 3, \beta_0 = 0.5, \mu = 0.2, \lambda = 0.1, \bar{s} = 10, q_0 = 5, w = 0.25, b = 1, a = 0.00004.$$

The equilibrium distribution capital stock is depicted in Figure 1.




FIGURE 1. Capital stock in equilibrium distribution for all ages,  $s \in [0, \bar{s}]$ , based on the parameter values  $\alpha = 3, \beta_0 = 0.5, \mu = 0.2, \lambda = 0.1, \bar{s} = 10, q_0 = 5, w = 0.25, b = 1, a = 0.00004$ .




FIGURE 2. Investment in equilibrium distribution for all ages,  $s \in [0, \bar{s}]$ , based on the parameter values  $\alpha = 3, \beta_0 = 0.5, \mu = 0.2, \lambda = 0.1, \bar{s} = 10, q_0 = 5, w = 0.25, b = 1, a = 0.00004$ .

We see that capital goods are non-monotonic with respect to age. To understand this, Figure 2 depicts equilibrium distribution investment behavior, where investment is given by

$$u_1^*(s) = \frac{1}{2\beta_0} \left( \int_s^{\bar{s}} e^{-(\lambda+\mu)(j-s)} (b - 2aQ^*) \alpha dj - q_0 e^{-ws} \right),$$

with  $Q^*$  being the production quantity in the equilibrium distribution.

Acquiring capital goods of older age is more attractive because they are cheaper (see (6.1)). On the other hand their lifetime is shorter, so they generate less revenue, which make older capital goods less attractive. Figure 2 shows that the last effect dominates for the older ages. The first effect plays a major role for younger ages. This makes sense because the convexity of the unit cost of acquisition with respect to age, as expressed in (6.1), makes that these capital goods get cheaper very quickly for slightly older age.

At first sight it is strange that  $K^*(\bar{s}) > 0$ , because  $\bar{s}$  is the age capital goods are scrapped. However, the presence of the adjustment costs,  $\frac{1}{2}\beta_0 [u_1(t, s)]^2$ , makes that it is not optimal

to sell all capital goods of age  $\bar{s}$ . In fact, convex adjustment costs make investments continuous over time, and thus also over age since age and time go together. Therefore, some of the capital goods of older age are still left. This is confirmed in the investment graph of Figure 2, where we also see that  $I^*(\bar{s})$  is negative.

If we leave out the effect that older capital goods are less costly, the effect of having a shorter lifetime when capital goods get older remains, and steady state investments decrease with age. This holds when we put

$$(6.10) \quad q_0 = q_1(s) = 0,$$

and, combining this with (6.4), (6.5), and (6.6), we obtain, when revenue is not specified, that

$$(6.11) \quad \begin{aligned} K^*(s) &= \eta w_1(s) \\ &= \frac{\eta\alpha}{2\beta_0(\mu+\lambda)} \left( e^{-\mu s} - e^{-\mu s} e^{-(\mu+\lambda)\bar{s}} + \frac{1}{\mu} (1 - e^{-\mu s}) \right) \\ &\quad - \frac{\eta\alpha}{2\beta_0(\mu+\lambda)} \left( e^{-(\mu+\lambda)\bar{s}} \frac{1}{2\mu+\lambda} (e^{(\mu+\lambda)s} - e^{-\mu s}) \right). \end{aligned}$$

In the specific case of a quadratic revenue function we get

$$(6.12) \quad \begin{aligned} K^*(s) &= \frac{bw_1(s)}{1+2ac_1} \\ &= \frac{b \left( \left( e^{-\mu s} - e^{-\mu s} e^{-(\mu+\lambda)\bar{s}} + \frac{1}{\mu} (1 - e^{-\mu s}) - e^{-(\mu+\lambda)\bar{s}} \frac{1}{2\mu+\lambda} (e^{(\mu+\lambda)s} - e^{-\mu s}) \right) \right)}{\frac{2\beta_0(\mu+\lambda)}{\alpha} + 2a\alpha \left( \frac{1-\mu}{\mu^2} (e^{-\mu\bar{s}} - 1) + \frac{2\mu+\lambda-1}{\mu(2\mu+\lambda)} e^{-(2\mu+\lambda)\bar{s}} + \frac{1}{\mu}\bar{s} - \frac{1}{(2\mu+\lambda)(\mu+\lambda)} + \frac{1-\mu-\lambda}{\mu(\mu+\lambda)} e^{-(\mu+\lambda)\bar{s}} \right)}. \end{aligned}$$

Equilibrium distribution investments being decreasing with age, also result in a hump-shaped structure of the steady state capital stock, like in Figure 1. The following proposition proves this analytically for a general revenue function, thus based on the equilibrium distribution capital stock specified in (6.11).

**Proposition 6.1.** *Consider the vintage capital stock model (2.1), (2.2), ((6.2)-(6.3)) with purely quadratic investment costs, i.e. we have (6.10) that partly replaces (6.1). Then equilibrium distribution capital stock  $K^*(s)$  is positive for all  $s \in [0, \bar{s}]$ , is increasing in age for  $s \in [0, s^*]$  and decreasing in age for  $s \in [s^*, \bar{s}]$ , where*

$$(6.13) \quad s^* = \frac{1}{2\mu+\lambda} \ln \left( \frac{2\mu+\lambda}{\mu+\lambda} \left( (1-\mu) e^{(\mu+\lambda)\bar{s}} + \mu \left( 1 - \frac{1}{2\mu+\lambda} \right) \right) \right) > 0.$$

Furthermore, it holds that

$$(6.14) \quad K^*(0) = \frac{\eta v}{2\beta_0(\mu+\lambda)} (1 - e^{-(\mu+\lambda)\bar{s}}) > 0,$$

(6.15)

$$K^*(\bar{s}) = \frac{\eta\alpha}{2\beta_0(\mu + \lambda)} \left( -e^{-\mu\bar{s}} \left( -1 + \frac{1}{\mu} \right) + e^{-\mu\bar{s}} e^{-(\mu+\lambda)\bar{s}} \left( \frac{1}{2\mu + \lambda} - 1 \right) + \frac{1}{\mu} - \frac{1}{2\mu + \lambda} \right) > 0.$$

*Proof.* From (6.12) we obtain that

$$K^{*'}(s) = \frac{\eta\alpha}{2\beta_0(\mu + \lambda)} e^{-\mu s} \left( \left( 1 - \mu + \mu \left( 1 - \frac{1}{2\mu + \lambda} \right) e^{-(\mu+\lambda)\bar{s}} - \frac{\mu + \lambda}{2\mu + \lambda} e^{-(\mu+\lambda)(\bar{s}-s)} e^{\mu s} \right) \right),$$

from which it is straightforwardly concluded that  $K^{*'}(s) > 0$  for  $s < s^*$ , with  $s^*$  given by (6.13) and vice versa.

To check whether  $s^*$  is positive we need to show that

$$\frac{2\mu + \lambda}{\mu + \lambda} \left( (1 - \mu) e^{(\mu+\lambda)\bar{s}} + \mu \left( 1 - \frac{1}{2\mu + \lambda} \right) \right) > 1,$$

which holds because

$$\begin{aligned} & \frac{2\mu + \lambda}{\mu + \lambda} \left( (1 - \mu) e^{(\mu+\lambda)\bar{s}} + \mu \left( 1 - \frac{1}{2\mu + \lambda} \right) \right) \\ & > \frac{2\mu + \lambda}{\mu + \lambda} \left( (1 - \mu) + \mu \left( 1 - \frac{1}{2\mu + \lambda} \right) \right) = 1. \end{aligned}$$

From (6.11) we straightforwardly obtain the expressions (6.14) and (6.15). To prove that  $K^*(\bar{s}) > 0$  we have to show that

$$-e^{-\mu\bar{s}} \left( -1 + \frac{1}{\mu} \right) + e^{-\mu\bar{s}} e^{-(\mu+\lambda)\bar{s}} \left( \frac{1}{2\mu + \lambda} - 1 \right) + \frac{1}{\mu} - \frac{1}{2\mu + \lambda} > 0,$$

which is true since

$$\begin{aligned} & -e^{-\mu\bar{s}} \left( -1 + \frac{1}{\mu} \right) + e^{-\mu\bar{s}} e^{-(\mu+\lambda)\bar{s}} \left( \frac{1}{2\mu + \lambda} - 1 \right) + \frac{1}{\mu} - \frac{1}{2\mu + \lambda} \\ & > 1 - \frac{1}{\mu} + \frac{1}{2\mu + \lambda} - 1 + \frac{1}{\mu} - \frac{1}{2\mu + \lambda} = 0. \end{aligned}$$

□

As a final illustration of the interesting economic results that can be obtained, let us focus on the impact of the productivity parameter  $\alpha$ . Let us increase  $\alpha$  from its original value 3, as in (6.9), to  $\alpha = 12$ . The resulting equilibrium distribution capital stock is depicted in Figure 3. The equilibrium distribution capital stock is still hump-shaped, as in the previous figures, but the difference is that the firm buys more capital goods. Higher productivity makes investing in capital goods more worthwhile. If we increase the productivity parameter  $\alpha$  even further to  $\alpha = 24$ , we obtain a equilibrium distribution capital stock being depicted in Figure 4. Now the capital stock is smaller for all ages. Concavity of the revenue function results in some bounded optimal quantity level, which, due to the increased productivity, can be produced by less capital goods.

The non-monotonic behavior of the capital stock that is obtained when productivity parameter  $\alpha$  goes up, is an interesting result, from which an expected outcome of including




FIGURE 3. Capital stock in equilibrium distribution for every age  $s \in [0, \bar{s}]$  for  $\alpha = 12$  with remaining parameter values  $\beta_0 = 0.5, \mu = 0.2, \lambda = 0.1, \bar{s} = 10, q_0 = 5, w = 0.25, b = 1, a = 0.00004$ .




FIGURE 4. Capital stock in equilibrium distribution for all ages  $s \in [0, \bar{s}]$  for productivity parameter  $\alpha = 24$  with remaining parameter values  $\beta_0 = 0.5, \mu = 0.2, \lambda = 0.1, \bar{s} = 10, q_0 = 5, w = 0.25, b = 1, a = 0.00004$ .

technological progress in the form of process innovation can be predicted. Increased productivity first results in more investments, but when productivity increases even further, investments go down because the optimal quantity in this market can be produced by less capital stock. The latter feature is new, and was for instance not derived in Feichtinger et al. (2006).

The non-monotonicity dependence of the equilibrium distribution capital stock on the productivity parameter can also be analytically proved in the special case of purely quadratic investment costs, as we do in the next proposition.

**Proposition 6.2.** *In case of quadratic revenue (see (6.3)) and purely quadratic investment costs, the equilibrium distribution capital stock,  $K^*(s)$  is increasing with the productivity parameter  $\alpha$  for  $\alpha \in [0, \hat{\alpha}]$  and decreasing with  $\alpha$  for  $\alpha \in [\hat{\alpha}, \infty)$ , where*

$$\hat{\alpha} = \frac{1}{c_1 b} \left( \sqrt{c_2 + \frac{c_1 b^2}{2a}} - c_2 \right) > 0,$$

in which

$$(6.16) \quad c_1 = \frac{1}{2\beta_0(\mu + \lambda)} \left( \frac{1 - \mu}{\mu^2} (e^{-\mu\bar{s}} - 1) + \frac{2\mu + \lambda - 1}{\mu(2\mu + \lambda)} e^{-(2\mu + \lambda)\bar{s}} \right) \\ + \frac{1}{2\beta_0(\mu + \lambda)} \left( \frac{\bar{s}}{\mu} - \frac{1}{(2\mu + \lambda)(\mu + \lambda)} + \frac{1 - (\mu + \lambda)}{\mu(\mu + \lambda)} e^{-(\mu + \lambda)\bar{s}} \right),$$

$$(6.17) \quad c_2 = \frac{q_0}{2\beta_0(\mu - w)} \left( \frac{\mu - w - 1}{\mu} (1 - e^{-\mu\bar{s}}) + \frac{1}{w} (1 - e^{-w\bar{s}}) \right)$$

*Proof.* From (6.4)-(6.8), (6.16) and (6.17) we obtain that

$$\frac{\partial K^*}{\partial \alpha} = \frac{\partial}{\partial \alpha} \left( \frac{b\alpha - 2ac_2\alpha^2}{1 + 2ac_1\alpha^2} \right) \varphi_1(s),$$

in which

$$\varphi_1(s) = \frac{1}{2\beta_0(\mu + \lambda)} \left( e^{-\mu s} - e^{-\mu s} e^{-(\mu + \lambda)\bar{s}} + \frac{1}{\mu} (1 - e^{-\mu s}) - e^{-(\mu + \lambda)\bar{s}} \frac{1}{2\mu + \lambda} (e^{(\mu + \lambda)s} - e^{-\mu s}) \right) \\ > 0.$$

It follows that

$$\frac{\partial K^*}{\partial \alpha} = \frac{-2ac_1b\alpha^2 - 4ac_2\alpha + b}{(1 + 2ac_1\alpha^2)^2} \varphi_1(s).$$

Recognizing that the concave second order polynomial

$$-2ac_1b\alpha^2 - 4ac_2\alpha + b$$

has a negative root and a positive root being equal to  $\hat{\alpha}$ , gives the result of the proposition.  $\square$

**Remark 6.3.** Concerning the stability of the equilibrium distribution  $K^*$ , we observe that Remark 5.12 applies here and may imply, depending on the value of the parameters, that  $K^*$  is locally, or even globally, stable.

**6.2. Power Revenues.** In this section we derive that the same result as that in Section 6.1, i.e. equilibrium distribution capital stock is hump-shaped in  $\alpha$ , can be established for an alternative revenue function based on the iso-elastic inverse demand function

$$p = bQ^{-\frac{1}{\varepsilon}},$$

in which  $\varepsilon > 1$  is the demand elasticity. Then the revenue function is

$$R(Q) = bQ^\gamma$$

with  $\gamma = 1 - 1/\epsilon$ . Since this revenue function has infinite derivative for  $Q = 0$ , it is not a  $C^1$  function. Therefore, instead we employ the revenue function

$$(6.18) \quad R(Q) = b((\theta + Q)^\gamma - \theta),$$

which approximates  $R(Q) = bQ^\gamma$  for  $\theta$  small. We obtain from Lemma 5.10 (iii) that for the revenue function as defined in (6.18) and the investment costs being purely quadratic, which implies that  $c_2 = 0$ , the  $\eta$  from (6.4) is implicitly determined by

$$(6.19) \quad \eta(\theta + \eta c_1)^{1-\gamma} - b\gamma = 0.$$

To establish the effect of the productivity parameter  $\alpha$  on  $K^*(s)$  in the case of iso-elastic demand, we first determine how  $\eta$  depends on  $\alpha$ . To do so, we first obtain from (6.7) that  $c_1$  is a quadratic function of  $\alpha$ :

$$c_1 = f\alpha^2$$

with

$$\begin{aligned} f &= \frac{1}{2\beta_0(\mu + \lambda)} \left( \frac{1 - \mu}{\mu^2} (e^{-\mu\bar{s}} - 1) + \frac{2\mu + \lambda - 1}{\mu(2\mu + \lambda)} e^{-(2\mu + \lambda)\bar{s}} \right) \\ &\quad + \frac{1}{2\beta_0(\mu + \lambda)} \left( \frac{1}{\mu\bar{s}} - \frac{1}{(2\mu + \lambda)(\mu + \lambda)} + \frac{1 - (\mu + \lambda)}{\mu(\mu + \lambda)} e^{-(\mu + \lambda)\bar{s}} \right) \\ &> 0. \end{aligned}$$

This implies that we can rewrite (6.19) into

$$(6.20) \quad \eta(\theta + \eta f\alpha^2)^{1-\gamma} - b\gamma = 0.$$

From the implicit function theorem we obtain that

$$\frac{\partial \eta}{\partial \alpha} = -\frac{2(1 - \gamma) f \eta \alpha}{\eta(1 - \gamma) + f\alpha^2 \eta + \theta} < 0,$$

whereas we also conclude from (6.20) that

$$(6.21) \quad \lim_{\alpha \rightarrow \infty} \eta(\alpha) = 0.$$

Now we are ready to establish how  $K^*(s)$  depends on  $\alpha$ . From (6.5) and (6.11) we get

$$K^*(s) = \eta\alpha\varphi(s),$$

with

$$\varphi(s) = \frac{1}{2\beta_0(\mu + \lambda)} \left( (1 - e^{-(\mu + \lambda)\bar{s}}) e^{-\mu s} + \frac{1 - e^{-\mu s}}{\mu} - \frac{e^{-(\mu + \lambda)\bar{s}}}{2\mu + \lambda} (e^{(\mu + \lambda)s} - e^{-\mu s}) \right) > 0.$$

Hence, we obtain

$$\begin{aligned} \frac{\partial K^*(s)}{\partial \alpha} &= \varphi(s) \left( \frac{\partial \eta}{\partial \alpha} \alpha + \eta \right) \\ &= \varphi(s) \eta \left( -\frac{2(1-\gamma)f\alpha^2}{\eta(1-\gamma) + f\alpha^2\eta + \theta} + 1 \right) \\ &= \varphi(s) \eta \left( -\frac{2(1-\gamma)f}{\frac{\eta}{\alpha^2}(1-\gamma) + f\eta + \frac{\theta}{\alpha^2}} + 1 \right). \end{aligned}$$

Since  $\frac{\eta}{\alpha^2}(1-\gamma) + f\eta + \frac{\theta}{\alpha^2}$  is decreasing in  $\alpha$ , and, due to (6.21), it also holds that

$$\begin{aligned} -\frac{2(1-\gamma)f}{\frac{\eta}{\alpha^2}(1-\gamma) + f\eta + \frac{\theta}{\alpha^2}} + 1 &> 0 \text{ for } \alpha = 0, \\ -\frac{2(1-\gamma)f}{\frac{\eta}{\alpha^2}(1-\gamma) + f\eta + \frac{\theta}{\alpha^2}} + 1 &< 0 \text{ for } \alpha \rightarrow \infty, \end{aligned}$$

we can conclude that we have proved the following proposition.

**Proposition 6.4.** *In case of iso-elastic demand and purely quadratic investment costs, the equilibrium distribution capital stock,  $K^*(s)$ , is increasing with the productivity parameter  $\alpha$  for  $\alpha \in [0, \hat{\alpha}]$ , and decreasing with  $\alpha$  for  $\alpha \in [\hat{\alpha}, \infty)$ , where  $\hat{\alpha}$  is implicitly given by*

$$-\frac{2(1-\gamma)f\hat{\alpha}^2}{\eta(\hat{\alpha})(1-\gamma) + f\eta(\hat{\alpha}) + \theta} + 1 = 0,$$

with

$$\begin{aligned} f &= \frac{1}{2\beta_0(\mu + \lambda)} \left( \frac{1-\mu}{\mu^2} (e^{-\mu\bar{s}} - 1) + \frac{2\mu + \lambda - 1}{\mu(2\mu + \lambda)} e^{-(2\mu + \lambda)\bar{s}} + \frac{1}{\mu}\bar{s} \right) \\ &\quad - \frac{1}{2\beta_0(\mu + \lambda)} \left( \frac{1}{(2\mu + \lambda)(\mu + \lambda)} + \frac{1 - (\mu + \lambda)}{\mu(\mu + \lambda)} e^{-(\mu + \lambda)\bar{s}} \right) \\ &> 0. \end{aligned}$$

**Remark 6.5.** Concerning the stability of the equilibrium distribution  $K^*$ , we observe that Lemma 5.11 applies here and may imply, depending on the value of the parameters, that  $K^*$  is locally, or even globally, stable.

#### APPENDIX A. PROOFS OF SUBSECTION 4.2

We here present a detailed description of the material in Subsection 4.2 as well as all the proofs of the results there stated. Firstly we note that solutions of the ODEs in (4.6) have to be intended in mild form. That is expressed in the following definitions.

**Definition A.1.** *Let Assumptions 3.2 be satisfied. Let  $t \geq 0$  and let  $u \in L_\lambda^p(t, +\infty; U)$ . The mild solution of (3.1) is the function  $y \in L_{loc}^1(t, +\infty; V')$  given by*

$$(A.1) \quad y(\tau) = e^{(\tau-t)A}x + \int_t^\tau e^{(\tau-\sigma)A}Bu(\sigma)d\sigma, \quad \tau \in [t, +\infty[.$$

The mild solution of (4.4)-(4.5) is the function  $\pi : [t, +\infty[ \rightarrow V$  given by

$$(A.2) \quad \pi(\tau) = \int_{\tau}^{+\infty} e^{(A_1^* - \lambda)(\sigma - \tau)} g'_0(y(\sigma)) d\sigma.$$

A mild solution to the closed loop equation (4.3) is a function  $y \in L^1_{loc}(t, +\infty; V')$  satisfying

$$(A.3) \quad y(\tau) = e^{(\tau - t)A} x + \int_t^{\tau} e^{(\tau - \sigma)A} B(h_0^*)'(-B^* \Psi'(y(s))) d\sigma, \quad \tau \in [t, +\infty[.$$

**Lemma A.2.** *Let Assumption 3.2 hold, assume  $p \geq 2$ ,  $q = \frac{p}{p-1}$  and  $\lambda > (2\omega) \vee \omega$ . Let also  $u \in L^p_{\lambda}(t, +\infty; U)$ . Then:*

- (i)  $\pi$  given by (A.2) is well defined and belongs to  $C^0([t, +\infty[; V)$ ;
- (ii) if  $p > 2$  and  $\lambda > 0$ , then  $\pi \in L^q_{\lambda}(t, +\infty; V)$ ;
- (iii) if  $p = 2$  and  $\lambda > 0$ , then  $\pi \in L^2_{\lambda + \varepsilon}(t, +\infty; V) \cap L^2(t, T; V)$ ,  $\forall T < +\infty$ ,  $\varepsilon > 0$ ;
- (iv) if  $p = 2$ ,  $\lambda > 0$  and  $\omega < 0$ , then  $\pi \in L^2_{\lambda}(t, +\infty; V)$ .

*Proof.* We first prove (i). Note that by assumptions on  $g'_0$ , the integrand in (A.2) can be estimated as follows

$$|e^{(A_1^* - \lambda)(\sigma - \tau)} g'_0(y(\sigma))|_V \leq |g'_0|_{\mathcal{B}_1} e^{-(\lambda - \omega)(\sigma - \tau)} (1 + |y(\sigma)|_{V'}).$$

Since  $\lambda > \omega$ , to prove the first assertion, i.e. that  $\pi(\cdot)$  is well defined, it is enough to show that, for every  $\tau \geq t$ , the map  $\sigma \mapsto e^{-(\lambda - \omega)(\sigma - \tau)} |y(\sigma)|_{V'}$  is in  $L^1(\tau, +\infty)$ .

By Assumption 3.2 (see also [44, Lemma 4.5]) one has

$$(A.4) \quad \begin{aligned} |y(\sigma)|_{V'} &\leq e^{\omega(\sigma - t)} |x|_{V'} + \int_t^{\sigma} e^{\omega(\sigma - r)} \|B\|_{\mathcal{L}(U, V')} |u(r)|_U dr \\ &\leq C e^{\omega\sigma} \left( |x|_{V'} + \int_t^{\sigma} e^{-\omega r} |u(r)|_U dr \right) \\ &\leq C_1 e^{\omega\sigma} (1 + \rho(t, \sigma)^{\frac{1}{q}}) \end{aligned}$$

for suitable constants  $C$  (depending only on  $t$ ) and  $C_1$  (depending on  $t$ ,  $x$ , and  $u$ ), where

$$\rho(t, \sigma) = \begin{cases} |e^{q(\frac{\lambda}{p} - \omega)t} - e^{q(\frac{\lambda}{p} - \omega)\sigma}| & \lambda \neq \omega p \\ |t - \sigma| & \lambda = \omega p. \end{cases}$$

Hence

$$e^{-(\lambda - \omega)\sigma} |y(\sigma)|_{V'} \leq C_1 e^{-(\lambda - 2\omega)\sigma} (1 + \rho(t, \sigma)^{\frac{1}{q}}),$$

so that in the case  $\lambda \neq \omega p$  one obtains

$$(A.5) \quad e^{-(\lambda - 2\omega)\sigma} \rho(t, \sigma)^{\frac{1}{q}} = |e^{-q(\lambda - 2\omega)\sigma} e^{q(\frac{\lambda}{p} - \omega)t} - e^{-(\lambda - q\omega)\sigma}|^{\frac{1}{q}} \leq C_2 \left[ e^{-(\lambda - 2\omega)\sigma} \vee e^{-\frac{1}{q}(\lambda - q\omega)\sigma} \right]$$

for a suitable constant  $C_2$ , whereas in the case  $\lambda = \omega p$  one has

$$(A.6) \quad e^{-(\lambda - 2\omega)\sigma} \rho(t, \sigma)^{\frac{1}{q}} = e^{-(\lambda - 2\omega)\sigma} |t - \sigma|^{\frac{1}{q}}.$$

Since  $\lambda > (2\omega) \vee \omega$  then also  $\lambda > q\omega$  since  $q = p/(p-1) \in (1, 2]$ . Hence, for each  $\tau \geq t$ , the integrand in (A.2) is in  $L^1(\tau, +\infty; V)$ .

The proof that  $\pi \in C^0([t, +\infty); V)$  follows from the dominated convergence theorem and the fact that the above estimates does not depend on  $\tau$ , when  $\tau$  is taken in any bounded interval.

Now we prove (ii). We start by showing that  $p > 2$  implies  $\pi \in L_\lambda^q(t, +\infty; V)$ . From the estimates above, one has

$$(A.7) \quad \begin{aligned} |\pi(\tau)|_V &\leq \frac{|g'_0|_{\mathcal{B}_1}}{\lambda - \omega} + \frac{|g'_0|_{\mathcal{B}_1} C_1 e^{\omega\tau}}{\lambda - 2\omega} + |g'_0|_{\mathcal{B}_1} C_1 e^{(\lambda - \omega)\tau} \int_\tau^{+\infty} e^{-(\lambda - 2\omega)\sigma} \rho(t, \sigma)^{\frac{1}{q}} d\sigma \\ &\equiv \gamma_1(\tau) + \gamma_2(\tau) + \gamma_3(\tau). \end{aligned}$$

The function  $\gamma_1$  is trivially in  $L_\lambda^q(t, +\infty; \mathbb{R})$ , since  $\lambda > 0$ . The function  $\gamma_2$  is in  $L_\lambda^q(t, +\infty; \mathbb{R})$  since, as observed above, it must be  $\lambda > q\omega$  as  $q \in (1, 2]$ . Regarding  $\gamma_3$ , in the case  $\lambda = \omega p$ , it must be necessarily  $\omega > 0$  (if not we cannot have  $\lambda > \omega$ ). Let then  $\delta > 0$  such that  $\lambda > 2\omega + 2\delta$ . Since, by simple computations,  $(\sigma - t)^{1/q} e^{-\delta\sigma} \leq \sigma^{1/q} e^{-\delta\sigma} \leq (qe\delta)^{-1/q}$ , we then have

$$\int_\tau^{+\infty} e^{-(\lambda - 2\omega)\sigma} |t - \sigma|^{\frac{1}{q}} d\sigma \leq (qe\delta)^{-1/q} \int_\tau^{+\infty} e^{-(\lambda - 2\omega - \delta)\sigma} d\sigma \leq (qe\delta)^{-1/q} \frac{e^{-(\lambda - 2\omega - \delta)\tau}}{\lambda - 2\omega - \delta}$$

Hence, in this case, for a suitable constant  $C_3$  one has

$$(A.8) \quad e^{-\lambda\tau} \gamma_3(t)^q \leq C_3 e^{-\lambda\tau} e^{q(\lambda - \omega)\tau} e^{-q(\lambda - 2\omega - \delta)\tau} = C_3 e^{-[\lambda - (\omega + \delta)q]\tau},$$

the last is an integrable function in  $[t, +\infty)$  by the choice of  $\delta$ . In the case  $\lambda \neq \omega p$ , by means of (A.5) one has

$$(A.9) \quad \int_\tau^{+\infty} e^{-(\lambda - 2\omega)\sigma} \rho(t, \sigma)^{\frac{1}{q}} d\sigma \leq C_2 \int_\tau^{+\infty} \left[ e^{-(\lambda - 2\omega)\sigma} \vee e^{-\frac{1}{q}(\lambda - q\omega)\sigma} \right] d\sigma.$$

By (A.7) we then have, for a suitable constant  $C_4 > 0$ ,

$$(A.10) \quad e^{-\lambda\tau} \gamma_3(\tau)^q \leq C_4 e^{-\lambda\tau} e^{q(\lambda - \omega)\tau} \left[ e^{-(\lambda - 2\omega)\tau} \vee e^{-\frac{1}{q}(\lambda - q\omega)\tau} \right]^q \leq C_4 \left[ e^{-(\lambda - q\omega)\tau} \vee e^{-\lambda(2 - q)\tau} \right],$$

which implies  $\gamma_3 \in L_\lambda^q(t, +\infty; \mathbb{R})$  also in this case.

To prove (iii) it suffices to observe that  $p = 2$  then (A.8)-(A.10) computed with  $q = 2$  and  $\lambda + \varepsilon$  in place of  $\lambda$  imply promptly  $\pi \in L_{\lambda + \varepsilon}^2(t, +\infty; V)$ ,  $\forall \varepsilon > 0$ .

Finally we prove (iv). Let  $p = 2$ . Observe that, for  $\tau \geq t$ ,

$$|\pi(\tau)|_V = \left| \int_\tau^{+\infty} e^{(A_1^* - \lambda)(\sigma - \tau)} g'_0(y(\sigma)) d\sigma \right|_V \leq |g'_0|_{\mathcal{B}_1} \int_\tau^{+\infty} e^{-(\lambda - \omega)(\sigma - \tau)} (1 + |y(\sigma)|_{V'}) d\sigma.$$

which implies

$$e^{-\lambda\tau} |\pi(\tau)|_V^2 \leq e^{-\lambda\tau} 2 |g'_0|_{\mathcal{B}_1}^2 \left[ \frac{1}{(\lambda - \omega)^2} + \left( \int_\tau^{+\infty} e^{-(\lambda - \omega)(\sigma - \tau)} |y(\sigma)|_{V'} d\sigma \right)^2 \right].$$

Since  $\lambda > 0$ , the first term is integrable on  $[t, +\infty[$ . Concerning the second term we exploit Jensen's inequality to get

$$(A.11) \quad \left( \int_{\tau}^{+\infty} e^{-(\lambda-\omega)(\sigma-\tau)} |y(\sigma)|_{V'}^2 d\sigma \right)^2 \leq \frac{1}{\lambda-\omega} \int_{\tau}^{+\infty} e^{-(\lambda-\omega)(\sigma-\tau)} |y(\sigma)|_{V'}^2 d\sigma$$

Hence

$$(A.12) \quad \int_t^{+\infty} e^{-\lambda\tau} |\pi(\tau)|_V^2 d\tau \leq \int_t^{+\infty} e^{-\lambda\tau} 2|g'_0|_{\mathcal{B}_1}^2 \left[ \frac{1}{(\lambda-\omega)^2} + \frac{1}{\lambda-\omega} \int_{\tau}^{+\infty} e^{-(\lambda-\omega)(\sigma-\tau)} |y(\sigma)|_{V'}^2 d\sigma \right] d\tau.$$

The first term of the right hand side is  $2|g'_0|_{\mathcal{B}_1}^2 e^{-\lambda t}/(\lambda-\omega)^2$ . To estimate the second we use Fubini-Tonelli Theorem (see e.g. [36, Theorem 1.33]), recalling that the integrand here is positive. Indeed

$$\begin{aligned} \int_t^{+\infty} e^{-\lambda\tau} \int_{\tau}^{+\infty} e^{-(\lambda-\omega)(\sigma-\tau)} |y(\sigma)|_{V'}^2 d\sigma d\tau &= \int_t^{+\infty} |y(\sigma)|_{V'}^2 e^{-(\lambda-\omega)\sigma} \int_t^{\sigma} e^{-\lambda\tau} e^{(\lambda-\omega)\tau} d\tau d\sigma \\ &= \int_t^{+\infty} |y(\sigma)|_{V'}^2 e^{-(\lambda-\omega)\sigma} \frac{e^{-\omega t} - e^{-\omega\sigma}}{\omega} d\sigma =: I_0 \end{aligned}$$

where, in the last step, we use that  $\omega \neq 0$ . To prove the claim it is now enough to prove that the last integral  $I_0$  is finite. First of all, by (A.4), we have

$$(A.13) \quad |y(\sigma)|_{V'}^2 \leq C_5 e^{2\omega\sigma} \left[ 1 + \left( \int_t^{\sigma} e^{-\omega r} |u(r)|_U dr \right)^2 \right]$$

for a suitable constant  $C_5$  (depending on  $t$ ,  $x$ , and  $u$ ). We apply again Jensen's inequality getting

$$\left( \int_t^{\sigma} e^{-\omega r} |u(r)|_U dr \right)^2 \leq \frac{e^{-\omega t} - e^{-\omega\sigma}}{\omega} \int_t^{\sigma} e^{-\omega r} |u(r)|_U^2 dr$$

Then we have

$$(A.14) \quad \begin{aligned} I_0 &\leq C_5 \left[ \int_t^{+\infty} e^{-(\lambda-3\omega)\sigma} \frac{e^{-\omega t} - e^{-\omega\sigma}}{\omega} d\sigma \right. \\ &\quad \left. + \int_t^{+\infty} e^{-(\lambda-3\omega)\sigma} \left( \frac{e^{-\omega t} - e^{-\omega\sigma}}{\omega} \right)^2 \int_t^{\sigma} e^{-\omega r} |u(r)|_U^2 dr d\sigma \right] \end{aligned}$$

Since  $\lambda > 0$  and  $\omega < 0$  the first integral of the right hand side of (A.14) is finite, positive, and, its value is

$$e^{-(\lambda-2\omega)t} \frac{1}{\omega} \left[ \frac{1}{\lambda-3\omega} - \frac{1}{\lambda-2\omega} \right] > 0.$$

Concerning the second integral of the right hand side of (A.14) (recalling that the integrand is positive) we apply Fubini-Tonelli Theorem again, to get that it is equal to

$$I_1 := \int_t^{+\infty} e^{-\omega r} |u(r)|_U^2 \int_r^{+\infty} e^{-(\lambda-3\omega)\sigma} \left( \frac{e^{-\omega t} - e^{-\omega\sigma}}{\omega} \right)^2 d\sigma dr$$

At this point we really need to use that  $\omega < 0$ <sup>8</sup>. This implies that the squared fraction above is smaller than  $e^{-2\omega\sigma}/(\omega^2)$ . Hence we have

$$I_1 \leq \frac{1}{\omega^2} \int_t^{+\infty} e^{-\omega r} |u(r)|_U^2 \int_r^{+\infty} e^{-(\lambda-\omega)\sigma} d\sigma dr \leq \frac{1}{\omega^2(\lambda-\omega)} \int_t^{+\infty} e^{-\lambda r} |u(r)|_U^2 dr$$

Since  $u \in L_\lambda^2(t, +\infty; U)$  the above imply the finiteness of  $I_1$  and, consequently, of  $I_0$ , which proves the claim.  $\square$

**Theorem A.3.** *Let Assumption 3.2 hold, let  $p \geq 2$ ,  $q = \frac{p}{p-1}$  and  $\lambda > (2\omega) \vee \omega$ . Let also  $u \in L_\lambda^p(t, +\infty; U)$ . If  $\pi \in W^{1,1}(t, +\infty; V)$  satisfies (4.4) almost everywhere in  $[t, +\infty)$  and the transversality condition (4.5), then  $\pi$  is given by (A.2), that is  $\pi$  is the mild solution of (4.4)-(4.5).*

*Proof.* By variation of constants formula, any  $\pi$  satisfying (4.4) a.e. must also satisfy

$$(A.15) \quad \pi(\tau) = e^{(A_1^* - \lambda)(T - \tau)} \pi(T) + \int_\tau^T e^{(A_1^* - \lambda)(\sigma - \tau)} g'_0(y(\sigma)) d\sigma, \quad \forall T > t, \quad \forall \tau \in [t, T].$$

Note that (4.5) implies

$$\lim_{T \rightarrow +\infty} |e^{(A_1^* - \lambda)(T - \tau)} \pi(T)|_V \leq \lim_{T \rightarrow +\infty} e^{(\omega - \lambda)(T - \tau)} |\pi(T)|_V = 0.$$

hence by passing to limits as  $T \rightarrow +\infty$  in (A.15) one derives

$$\pi(\tau) = \lim_{T \rightarrow +\infty} \int_\tau^T e^{(A_1^* - \lambda)(\sigma - \tau)} g'_0(y(\sigma)) d\sigma = \int_\tau^{+\infty} e^{(A_1^* - \lambda)(\sigma - \tau)} g'_0(y(\sigma)) d\sigma$$

where the last equality follows from estimates (A.5)-(A.6).  $\square$

We are now ready to prove the Maximum Principle.

### Proof of Theorem 4.5 (Maximum Principle)

Let  $K : L_\lambda^p(t, +\infty; U) \rightarrow \mathbb{R} \cup \{+\infty\}$ , and  $G : L_\lambda^p(t, +\infty; U) \rightarrow \mathbb{R} \cup \{+\infty\}$  be defined by

$$K(u) := \int_t^{+\infty} e^{-\lambda\tau} h_0(u(\tau)) d\tau, \quad G(u) := \int_t^{+\infty} e^{-\lambda\tau} g_0(y(\tau; t, x, u)) d\tau$$

so that for all  $u \in \text{dom}(K) \cap \text{dom}(G)$  we have

$$(A.16) \quad J(u) = K(u) + G(u), \quad \text{and} \quad \partial J(u) \supseteq \partial K(u) + \partial G(u).$$

*Claim 1:* For  $u \in \text{int dom}(K)$  we have

$$(A.17) \quad \partial K(u) = S, \quad \text{where} \\ S \equiv \{\varphi \in L_\lambda^q(t, +\infty; U) : \varphi(\tau) \in \partial h_0(u(\tau)), \text{ for a.e. } \tau \in [t, +\infty)\}$$

<sup>8</sup>This was not needed up to now. Above we only used the fact that  $\lambda > 3\omega$  and, to simplify computations,  $\omega \neq 0$ .

Indeed

(A.18)

$$\partial K(u) = \left\{ \varphi \in L_\lambda^q(t, +\infty; U) : \int_t^{+\infty} e^{-\lambda\tau} [h_0(w(\tau)) - h_0(u(\tau)) - (\varphi(\tau)|w(\tau) - u(\tau))_U] d\tau \geq 0, \forall w \in L_\lambda^p(t, +\infty; U) \right\},$$

so that  $S \subset \partial K(u)$  is straightforward. To show the reverse inclusion, we let  $\varphi$  be any fixed element of  $\partial K(u)$ ,  $E$  any measurable subset of  $[t, +\infty)$ , and we set, for any  $w \in L_\lambda^p(t, +\infty; U)$ ,

$$(A.19) \quad \tilde{w}(\tau) = \begin{cases} u(\tau), & \tau \notin E \\ w(\tau), & \tau \in E \end{cases}$$

Clearly we still have  $\tilde{w} \in L_\lambda^p(t, +\infty; U)$ , hence we derive

$$\int_E e^{-\lambda\tau} [h_0(w(\tau)) - h_0(u(\tau)) - (\varphi(\tau)|w(\tau) - u(\tau))_U] d\tau \geq 0, \forall w \in L_\lambda^p(t, +\infty; U).$$

Since  $E$  and  $w$  where arbitrarily chosen, the above implies

$$h_0(w(\tau)) - h_0(u(\tau)) - (\varphi(\tau)|w(\tau) - u(\tau))_U \geq 0, \text{ for a.e. } \tau \in [0, +\infty)$$

that is,  $\varphi(\tau) \in \partial h_0(\tau)$  for almost every  $\tau \geq 0$ .

*Claim 2:* Let  $(u, y)$  be admissible at  $(t, x)$ . Assume that there exists  $\pi \in L_\lambda^q(t, +\infty; V)$  such that  $(\pi, u, y)$  satisfies (4.6). Then  $(u, y)$  is optimal at  $(t, x)$ .

Let  $v$  be any control in  $\text{dom}(G)$ . Then

$$(A.20) \quad \begin{aligned} G(v) - G(u) &= \int_t^{+\infty} [g_0(y(\tau; v)) - g_0(y(\tau; u))] e^{-\lambda\tau} d\tau \\ &\geq \int_t^{+\infty} \langle g'_0(y(\tau; u)), \int_t^\tau e^{(\tau-\sigma)A} B(v(\sigma) - u(\sigma)) d\sigma \rangle e^{-\lambda\tau} d\tau \\ &= \int_t^{+\infty} \int_\sigma^{+\infty} (B^* e^{(\tau-\sigma)A_1^*} g'_0(y(\tau; u)) | v(\sigma) - u(\sigma))_U e^{-\lambda\tau} d\tau d\sigma \\ &= \int_t^{+\infty} \left( \int_\sigma^{+\infty} B^* e^{(\tau-\sigma)(A_1^* - \lambda)} g'_0(y(\tau; u)) d\tau | v(\sigma) - u(\sigma) \right)_U e^{-\lambda\sigma} d\sigma \\ &= \int_t^{+\infty} (B^* \pi(\sigma) | v(\sigma) - u(\sigma))_U e^{-\lambda\sigma} d\sigma \\ &= \langle B^* \pi, v - u \rangle_{L_\lambda^q(t, +\infty; U), L_\lambda^p(t, +\infty; U)} \end{aligned}$$

where we could exchange the order of integration since  $\pi$  is in  $L_\lambda^q(t, +\infty; V)$  by assumption. Then we proved that

$$B^* \pi \in \partial G(u).$$

Now, by (4.6) we also know that  $-B^*\pi(\sigma) \in \partial h_0(u(\sigma))$  almost everywhere in  $[t, +\infty[$ ; hence, by Claim 1, we get  $-B^*\pi \in \partial K(u)$ . By (A.16) it follows that  $\partial J(u) \ni 0$ , that is,  $u$  is optimal and (i) is proved.

*Claim 3.* Assume that, either  $p > 2$  and  $\lambda > 0$ , or  $p = 2$ ,  $\lambda > 0$  and  $\omega < 0$ . Assume that  $(u^*, y^*)$  is optimal at  $(t, x)$ , and let  $\pi^*$  be the associated solution of (A.2). Then  $G$  is Gâteaux differentiable in  $u^*$  with  $G'(u^*) = B^*\pi^*$ . Consequently  $\partial J(u^*) = B^*\pi^* + S$ .

First of all we recall that, by assumption and by Lemma A.2, we have  $\pi^* \in L_\lambda^q(t, +\infty; V)$ . Moreover, for any fixed  $v$  in  $L_\lambda^p(t, +\infty; U)$ , and any  $\epsilon > 0$ , there exists  $0 < \epsilon_0 \leq \epsilon$  such that

$$\begin{aligned} \frac{G(u^* + \epsilon v) - G(u^*)}{\epsilon} &= \int_t^{+\infty} \frac{g_0(y(\tau; u^* + \epsilon v)) - g_0(y^*(\tau))}{\epsilon} e^{-\lambda\tau} d\tau = \\ &= \int_t^{+\infty} \langle g'_0(y(\tau; u^* + \epsilon_0 v)), y(\tau; u^* + \epsilon_0 v) - y^*(\tau) \rangle e^{-\lambda\tau} d\tau \\ &= \int_t^{+\infty} \left\langle g'_0(y(\tau; u^* + \epsilon_0 v)), \int_t^\tau e^{(\tau-\sigma)A} Bv(\sigma) d\sigma \right\rangle e^{-\lambda\tau} d\tau \end{aligned}$$

Hence, arguing as in (A.20) to rewrite the term  $\langle B^*\pi^*, v \rangle_{L_\lambda^q, L_\lambda^p}$ , we get

$$\begin{aligned} &\left| \frac{G(u^* + \epsilon v) - G(u^*)}{\epsilon} - \langle B^*\pi^*, v \rangle_{L_\lambda^q, L_\lambda^p} \right| = \\ &= \left| \int_t^{+\infty} \frac{g_0(y(\tau; u^* + \epsilon v)) - g_0(y^*(\tau))}{\epsilon} e^{-\lambda\tau} d\tau - \langle B^*\pi^*, v \rangle_{L_\lambda^q, L_\lambda^p} \right| = \\ &= \left| \int_t^{+\infty} \langle g'_0(y(\tau; u^* + \epsilon_0 v)) - g'_0(y^*(\tau)), \int_t^\tau e^{(\tau-\sigma)A} Bv(\sigma) d\sigma \rangle e^{-\lambda\tau} d\tau \right| \\ &\leq [g'_0]_{\epsilon_0} \int_t^{+\infty} \left| \int_t^\tau e^{(\tau-\sigma)A} Bv(\sigma) d\sigma \right|_{V'}^2 e^{-\lambda\tau} d\tau =: I \end{aligned}$$

We estimate now the right hand side in the case when  $p > 2$  and  $\lambda \neq \omega p$ . By Hölder inequality one has

$$\begin{aligned} \left| \int_t^\tau e^{(\tau-\sigma)A} Bv(\sigma) d\sigma \right|_{V'} &\leq e^{\omega\tau} \left[ \int_t^\tau e^{(\frac{\lambda}{p}-\omega)q\sigma} d\sigma \right]^{\frac{1}{q}} \|B\|_{\mathcal{L}(U, V')} |v|_{L_\lambda^p(t, +\infty; U)} \\ &= e^{\omega\tau} \left| \frac{e^{(\frac{\lambda}{p}-\omega)q\tau} - e^{(\frac{\lambda}{p}-\omega)qt}}{q \left(\frac{\lambda}{p} - \omega\right)} \right|^{\frac{1}{q}} \|B\|_{\mathcal{L}(U, V')} |v|_{L_\lambda^p(t, +\infty; U)} \end{aligned}$$

Then, for a suitable constant  $C_5$ , depending on  $g$ ,  $B$  and  $v$ , we have

$$\begin{aligned}
(A.21) \quad I &\leq [g'_0]\epsilon_0 \int_t^{+\infty} \left| e^{\omega\tau} \left| \frac{e^{(\frac{\lambda}{p}-\omega)q\tau} - e^{(\frac{\lambda}{p}-\omega)qt}}{q \left( \frac{\lambda}{p} - \omega \right)} \right|^{\frac{1}{q}} \|B\|_{\mathcal{L}(U,V')} |v|_{L_\lambda^p(t,+\infty;U)} \right|^2 e^{-\lambda\tau} d\tau \\
&\leq C_5 \epsilon_0 \int_t^{+\infty} \left| e^{(\frac{\lambda}{p}-\omega)q\tau} - e^{(\frac{\lambda}{p}-\omega)qt} \right|^{\frac{2}{q}} e^{-(\lambda-2\omega)\tau} d\tau \\
&\leq C_5 \epsilon_0 \int_t^{+\infty} \left( e^{(\frac{\lambda}{p}-\omega)q\tau} \vee 1 \right)^{\frac{2}{q}} e^{-(\lambda-2\omega)\tau} d\tau \\
&= C_5 \epsilon_0 \int_t^{+\infty} e^{\lambda(\frac{2}{p}-1)\tau} \vee e^{-(\lambda-2\omega)\tau} d\tau.
\end{aligned}$$

Since  $p > 2 \Rightarrow \lambda \left( \frac{2}{p} - 1 \right) < 0$ , then one may let  $\epsilon \rightarrow 0$  and obtains that the right hand side in (A.21) goes to 0.

Let now  $p > 2$  and  $\lambda = \omega p$ . By Hölder inequality one has

$$\begin{aligned}
\left| \int_t^\tau e^{(\tau-\sigma)A} Bv(\sigma) d\sigma \right|_{V'} &\leq \|B\|_{\mathcal{L}(U,V')} e^{\omega\tau} \int_t^\tau e^{-\frac{\lambda}{p}\sigma} |v(\sigma)|_U d\sigma \\
&\leq \|B\|_{\mathcal{L}(U,V')} e^{\omega\tau} |\tau - t|^{\frac{1}{q}} |v|_{L_\lambda^p(t,+\infty;U)}
\end{aligned}$$

Then, for suitable  $C_6 > 0$  we get

$$\begin{aligned}
I &\leq [g'_0]\epsilon_0 \int_t^{+\infty} \left| \|B\|_{\mathcal{L}(U,V')} e^{\omega\tau} |\tau - t|^{\frac{1}{q}} |v|_{L_\lambda^p(t,+\infty;U)} \right|^2 v^2 e^{-\lambda\tau} d\tau \\
&\leq C_6 \epsilon_0 \int_t^{+\infty} |\tau - t|^{\frac{2}{q}} e^{-(\lambda-2\omega)\tau} d\tau
\end{aligned}$$

and by letting  $\epsilon \rightarrow 0$  the right hand side goes to 0.

Let now  $p = 2$ ,  $\lambda > 0$  and  $\omega < 0$ . Then we have

$$\begin{aligned}
\left| \int_t^\tau e^{(\tau-\sigma)A} Bv(\sigma) d\sigma \right|_{V'}^2 &\leq \|B\|_{\mathcal{L}(U,V')}^2 \left[ \int_t^\tau e^{\omega(\tau-\sigma)} |v(\sigma)| d\sigma \right]^2 \\
&\leq \|B\|_{\mathcal{L}(U,V')}^2 \frac{1}{\omega} [e^{\omega(\tau-t)} - 1] \int_t^\tau e^{\omega(\tau-\sigma)} |v(\sigma)|^2 d\sigma
\end{aligned}$$

where in the last step we used the Jensen's inequality. Hence, by using Fubini-Tonelli Theorem, we get

$$\begin{aligned}
& \int_t^{+\infty} \left| \int_t^\tau e^{(\tau-\sigma)A} Bv(\sigma) d\sigma \right|_{V'}^2 e^{-\lambda\tau} d\tau \leq \\
& \leq \|B\|_{\mathcal{L}(U,V')}^2 \int_t^{+\infty} \frac{1}{\omega} [e^{\omega(\tau-t)} - 1] e^{-\lambda\tau} \int_t^\tau e^{\omega(\tau-\sigma)} |v(\sigma)|^2 d\sigma d\tau = \\
& = \|B\|_{\mathcal{L}(U,V')}^2 e^{-\lambda t} \int_t^{+\infty} e^{\omega(\sigma-t)} |v(\sigma)|^2 \int_\sigma^{+\infty} \frac{1}{\omega} [e^{\omega(\tau-t)} - 1] e^{-(\lambda-\omega)(\tau-t)} d\tau d\sigma = \\
& = \|B\|_{\mathcal{L}(U,V')}^2 e^{-\lambda t} \int_t^{+\infty} |v(\sigma)|^2 \frac{1}{\omega} \left[ \frac{e^{-(\lambda-\omega)(\sigma-t)}}{\lambda-2\omega} - \frac{e^{-\lambda(\sigma-t)}}{\lambda-\omega} \right] d\sigma \leq \\
& \leq \|B\|_{\mathcal{L}(U,V')}^2 \frac{1}{(-\omega)(\lambda-\omega)} \int_t^{+\infty} |v(\sigma)|^2 e^{-\lambda\sigma} d\sigma
\end{aligned}$$

where, in the last inequality, we used that  $\omega < 0$ . This immediately implies that

$$I \leq [g'_0] \epsilon_0 \|B\|_{\mathcal{L}(U,V')}^2 \frac{1}{(-\omega)(\lambda-\omega)} |v|_{L^2_\lambda(t,+\infty;U)}$$

which immediately gives the claim.

*Claim 4.* Assume that, either  $p > 2$  and  $\lambda > 0$ , or  $p = 2$ ,  $\lambda > 0$  and  $\omega < 0$ . Assume that  $(u^*, y^*)$  is optimal at  $(t, x)$ , and let  $\pi^*$  be the associated solution of (A.2). Then  $(\pi^*, u^*, y^*)$  is a mild solution of (4.6).

We only need to prove that the last line of (4.6). From optimality of  $u^*$  we have  $\partial J(u^*) \ni 0$ . Then Claim 1 and Claim 3 imply (4.7) and Claim 4 follows.  $\square$

**Proof of Theorem 4.6** Firstly we prove that  $\pi^*(t; t, x) \in \partial Z_0(x)$ . We recall that in Theorem 4.5 we showed that  $-B^* \pi^*(\tau; t, x) \in \partial h_0(u^*(\tau))$  almost everywhere in  $[0, +\infty)$ . Then, for all  $\bar{x} \in V'$ , and an associated control  $\bar{u}$ , optimal at  $(t, \bar{x})$ , we have

$$\begin{aligned}
Z_0(\bar{x}) - Z_0(x) &= e^{\lambda t} [J(t, \bar{x}, \bar{u}) - J(t, x, u^*)] \\
&\geq \int_t^{+\infty} \left[ \langle g'_0(y^*(\tau)), \bar{y}(\tau) - y^*(\tau) \rangle - (B^* \pi^*(\tau; t, x)) | \bar{u}(\tau) - u^*(\tau) \rangle_U \right] e^{-\lambda(\tau-t)} d\tau.
\end{aligned}$$

Note that

$$\begin{aligned}
& \int_t^{+\infty} \langle g'_0(y^*(\tau)), \bar{y}(\tau) - y^*(\tau) \rangle_{V,V'} e^{-\lambda(\tau-t)} d\tau = \\
& = \int_t^{+\infty} \left\langle g'_0(y^*(\tau)), e^{A(\tau-t)}(\bar{x} - x) + \int_t^\tau e^{A(\tau-\sigma)} B(\bar{u}(\sigma) - u^*(\sigma)) d\sigma \right\rangle e^{-\lambda(\tau-t)} d\tau = \\
& = \int_t^{+\infty} \langle e^{(A_1^* - \lambda)(\tau-t)} g'_0(y^*(\tau)), \bar{x} - x \rangle d\tau + \\
& \quad + \int_t^{+\infty} \int_t^\tau \langle g'_0(y^*(\tau)), e^{A(\tau-\sigma)} (B(\bar{u}(\sigma) - u^*(\sigma))) \rangle e^{-\lambda(\tau-t)} d\sigma d\tau.
\end{aligned}$$

The last term can be rewritten exchanging the integrals as

$$\begin{aligned} & \int_t^{+\infty} \int_\sigma^{+\infty} (B^* e^{(A_1^* - \lambda)(\tau - \sigma)} g'_0(y^*(\tau)) | \bar{u}(\sigma) - u^*(\sigma) )_U e^{-\lambda(\sigma - t)} d\tau d\sigma = \\ & = \int_t^{+\infty} (B^* \pi^*(\sigma; t, x) | \bar{u}(\sigma) - u^*(\sigma) )_U e^{-\lambda(\sigma - t)} d\sigma. \end{aligned}$$

Hence we get

$$Z_0(\bar{x}) - Z_0(x) \geq \left\langle \int_t^{+\infty} e^{(A_1^* - \lambda)(\tau - t)} g'_0(y^*(\tau)) d\tau, \bar{x} - x \right\rangle = \langle \pi^*(t; t, x), \bar{x} - x \rangle,$$

and the assertion is proven. The proof that  $\pi^*(\tau; \tau, y^*(\tau)) \in \partial Z_0(y^*(\tau))$  for every  $\tau \geq t$  is standard but we write it here for the sake of completeness. Let  $\tau > t$  and observe that, by the dynamic programming principle, the control defined by

$$u_{0, y^*(\tau)}(\sigma) \equiv u^*(\sigma + \tau)$$

is optimal at  $(0, y^*(\tau))$ . Consequently the associated trajectory satisfies

$$y(\sigma; 0, y^*(\tau), u_{0, y^*(\tau)}) = y(\sigma + \tau; \tau, y^*(\tau), u^*) = y^*(\sigma + \tau).$$

Then by the first part of the proof we have

$$\begin{aligned} \partial Z_0(y^*(\tau)) \ni \pi^*(0; 0, y^*(\tau)) &= \\ &= \int_0^{+\infty} e^{(A_1^* - \lambda)\sigma} g'_0(y(r; 0, y^*(\tau), u_{0, y^*(\tau)})) d\sigma \\ &= \int_0^{+\infty} e^{(A_1^* - \lambda)\sigma} g'_0(y^*(\sigma + \tau)) d\sigma \\ &= \int_\tau^{+\infty} e^{(A_1^* - \lambda)(r - \tau)} g'_0(y^*(r)) dr \\ &= \pi^*(\tau; \tau, y^*(\tau)) \end{aligned}$$

which gives the claim. It finally suffices to note that, when  $p > 2$ , one has by Theorem 4.2, that  $Z_0 = \Psi$  and  $\partial \Psi(x) = \{\Psi'(x)\}$  to complete the proof.  $\square$

## APPENDIX B. PROOFS OF SUBSECTION 4.3

We here work out the proofs of the results stated in Subsection 4.3.

**Proof of Theorem 4.10.** We first prove (i). Assume  $(\bar{x}, \bar{\pi}, \bar{u})$  is a MP-equilibrium point for problem (P). Since  $\lambda > \omega$ , then  $\lambda \in \rho(A_1^*)$ , and the second of (4.11) applies, implying

$$(B.1) \quad \bar{\pi} = (\lambda - A_1^*)^{-1} g'_0(\bar{x}).$$

Then, plugging (B.1) into the third equation of (4.9) and then  $\bar{u}$  so obtained into the first, we derive (4.13). Moreover, using (B.1) in Theorem 4.6, we get (4.14).

We now prove (ii). We consider a CLE-equilibrium point  $\hat{x}$  and set  $\hat{u} := (h_0^*)'(-B^* \Psi'_0(\hat{x}))$ . By Theorem 4.3 and Remark 4.4 we know that the couple  $(\hat{x}, \hat{u})$  is optimal.

By Theorem 4.5-(ii) we can associate, to such optimal couple, a costate  $\hat{\pi}$  which is the (mild) solution of the costate equation in (4.6). Such mild solution is then necessarily stationary (see Definition A.1) and, since  $\lambda > \omega$ , given by

$$\hat{\pi} \equiv (\lambda - A_0^*)^{-1} g'_0(\hat{x}).$$

Moreover, by Theorem 4.6 it is also true that  $\hat{\pi} = \Psi'(\hat{x})$ . As a consequence,  $(\hat{x}, \hat{\pi}, \hat{u})$  solves (4.9) and is then a MP-equilibrium point.  $\square$

Before demonstrating Lemma 4.11 we need to state and prove the following result.

**Proposition B.1.** *Assume that  $0 \in \rho(A_1^*)$  (that is,  $(A_1^*)^{-1}$  is well defined and bounded in  $H$ ). Then  $A^{-1}$  has bounded inverse on  $V'$ , defined by the position*

$$\langle A^{-1}f, \varphi \rangle = \langle f, (A_1^*)^{-1}\varphi \rangle, \text{ for all } f \in V' \text{ and } \varphi \in V.$$

Moreover

$$\|A^{-1}\|_{\mathcal{L}(V')} \leq \|(A_1^*)^{-1}\|_{\mathcal{L}(H)}.$$

*Proof.* For all  $f \in V'$  and  $\varphi \in V$  we have

$$\begin{aligned} |\langle A^{-1}f, \varphi \rangle| &= |\langle f, (A_1^*)^{-1}\varphi \rangle| \\ &\leq |f|_{V'} |(A_1^*)^{-1}\varphi|_V \\ (B.2) \quad &= |f|_{V'} (|(A_1^*)^{-1}\varphi|_H + |A_1^*(A_1^*)^{-1}\varphi|_H) \\ &= |f|_{V'} (|(A_1^*)^{-1}\varphi|_H + |(A_1^*)^{-1}A_1^*\varphi|_H) \\ &\leq |f|_{V'} \|(A_1^*)^{-1}\|_{\mathcal{L}(H)} |\varphi|_V \end{aligned}$$

$\square$

**Proof of Lemma 4.11.**

Define  $T : V' \rightarrow V'$  as in (4.16). By its definition,  $T$  satisfies

$$(B.3) \quad |Tx - Ty|_{V'} \leq \|A^{-1}\|_{\mathcal{L}(V')} \|B\|_{\mathcal{L}(U, V')}^2 [(h_0^*)'] [g_0] \frac{1}{\lambda - \omega} |x - y|_{V'},$$

implying the claim. Since  $T(V') \subseteq D(A)$ , then fixed points of  $T$  lie in  $D(A)$ .  $\square$

**Proof of Proposition 4.16** Let  $x \in I(\bar{x})$  and let  $x^*(t)$  be the associated optimal trajectory, i.e. the solution of the closed loop equation

$$(B.4) \quad \begin{cases} y'(t) = Ay(t) + f(y(t)), & t > 0 \\ y(0) = x, \end{cases}$$

Let  $x_n \in I(\bar{x}) \cap D(A)$  be such that  $x_n \rightarrow x$  in  $V'$  as  $n \rightarrow +\infty$ . Let  $y_n^*(t)$  be the associated optimal trajectory. Since  $y_n^*$  is continuous, then it must remain in  $I(\bar{x})$  at least for a sufficiently small  $t$ . For such  $t$  we must have (recall that  $y_n^*(t) \in D(A)$  by assumption)

$$\begin{aligned} \frac{1}{2} \frac{d}{dt} |y_n^*(t) - \bar{x}|_{V'}^2 &= ((y_n^*(t))' |y_n^*(t) - \bar{x})_{V'} \\ &= (A(y_n^*(t) - \bar{x}) + f(y_n^*(t)) - f(\bar{x}) |y_n^*(t) - \bar{x})_{V'} \leq \xi |y_n^*(t) - \bar{x}|_{V'}^2. \end{aligned}$$

This implies that

$$(B.5) \quad |y_n^*(t) - \bar{x}|_{V'}^2 \leq e^{2\xi t} |x_n - \bar{x}|_{V'}^2$$

Next we take the limits as  $n \rightarrow +\infty$ . Note that  $z_n(t) := y_n^*(t) - x^*(t)$  solves

$$(B.6) \quad \begin{cases} z_n'(t) = Az_n(t) + f(y_n^*(t)) - f(x^*(t)), & t > 0 \\ z(0) = x_n - x. \end{cases}$$

By Theorem 4.3  $f$  is Lipschitz continuous so with Lipschitz constant  $[f]_{0,1}$ , a standard Gronwall inequality implies

$$|y_n^*(t) - x^*(t)|_{V'} \leq |x_n - x|_{V'} e^{(\omega + [f]_{0,1})t},$$

hence  $|y_n^*(t) - x^*(t)|_{V'}$  converges to 0 for every  $t \geq 0$ . Consequently, from (B.5) one has

$$|x^*(t) - \bar{x}|_{V'}^2 \leq e^{2\xi t} |x - \bar{x}|_{V'}^2, \quad \text{for every } t \geq 0,$$

implying the claims.  $\square$

**Proof of Corollary 4.17.** It is enough to check that (4.18) is verified, either in  $I$ , or in  $V'$ . If  $f$  is Lipschitz continuous on  $I$  (in the topology of  $V'$ ), with Lipschitz constant  $\bar{\theta} < \theta$  then we have

$$(f(x) - f(\bar{x}))|x - \bar{x}|_{V'} \leq \bar{\theta} |x - \bar{x}|_{V'}^2, \quad \forall x \in I.$$

This immediately implies that (4.18) holds in  $I$  with  $\xi = \bar{\theta} - \theta$ .

Similarly, if  $f$  is globally Lipschitz continuous in  $V'$ , we get that (4.18) is verified in the whole  $V'$ .  $\square$

## APPENDIX C. PROOFS OF SECTION 5

**Proof of Lemma 5.4.** Note that if  $A_0^*$  and  $A$  are the operators described in Section 3.3, then the following two facts hold true. First, by definition of  $A_0^*$ , we have  $\bar{\alpha} = (\lambda - A_0^*)^{-1}\alpha$ . Second,  $A$  is invertible, so that equation (4.13) may be rewritten as  $Tx = x$  by means of the operator  $T$  defined in (4.16). Now (3.9) holds, so that one has

$$g_0'(x) = -R'(\langle \alpha, x \rangle)\alpha, \quad \text{and} \quad (\lambda - A_0^*)^{-1}g_0'(x) = -R'(\langle \alpha, x \rangle)\bar{\alpha}$$

Moreover, since  $h_0(u) = C(u_0, u_1) = C_0(u_0) + C_1(u_1)$ , then the convex conjugate  $C^*$  of  $C$  is also of type  $C^*(u_0, u_1) = C_0^*(u_0) + C_1^*(u_1)$ . Then, recalling the definition of  $B, B^*$  in Section 2, and by means of (5.5), the operator  $T$  defined in (4.16) can be rewritten as

follows:

$$\begin{aligned}
Tx &= -A^{-1}B(C^*)'(B^*R'(\langle\alpha, x\rangle)\bar{\alpha}) \\
&= -A^{-1}B(C^*)'(R'(\langle\alpha, x\rangle)\bar{\alpha}(0), R'(\langle\alpha, x\rangle)\bar{\alpha}) \\
&= -A^{-1}B((C_0^*)'(R'(\langle\alpha, x\rangle)\bar{\alpha}(0)), (C_1^*)'(R'(\langle\alpha, x\rangle)\bar{\alpha})) \\
&= -A^{-1}[(C_0^*)'(R'(\langle\alpha, x\rangle)\bar{\alpha}(0))\delta_0 + (C_1^*)'(R'(\langle\alpha, x\rangle)\bar{\alpha})] \\
&= (C_0^*)'(R'(\langle\alpha, x\rangle)\bar{\alpha}(0))[-A^{-1}\delta_0] + [-A^{-1}](C_1^*)'(R'(\langle\alpha, x\rangle)\bar{\alpha})
\end{aligned}$$

which by (3.7) implies (5.6). By Theorem 4.10 we derive the remaining statements.  $\square$

**Proof of Theorem 5.5.** Set  $\eta : V' \rightarrow \mathbb{R}$ ,  $\eta(x) = R'(\langle\alpha, x\rangle)$  and note that, by (5.6), we get  $Tx(s) = F(\eta(x))(s)$ . Then the equation  $Tx = x$  is rewritten as

$$F(\eta(x))[s] = x(s), \quad \forall s \in [0, \bar{s}].$$

Applying  $\eta$  on both sides of such equation we get

$$R'(\langle\alpha, F(\eta(x))\rangle) = \eta(x).$$

Hence, if  $\bar{x}$  is a solution of  $Tx = x$ , then  $\eta(\bar{x})$  is a solution of (5.10). Viceversa, let  $\bar{\eta}$  be a solution to (5.10). Then, substituting into (5.8) and (5.6), we get that  $F(\bar{\eta})$  solves  $F(\bar{\eta}) = TF(\bar{\eta})$ , so  $F(\bar{\eta})$  solves  $Tx = x$ .  $\square$

**Proof of Theorem 5.11.** Note that when the value function  $V_0$  is differentiable, then Proposition 4.15 applies with  $\Psi = -V_0$ . Indeed by Lemma 3.5 and the Lumer-Philips Theorem (see e.g. [68, Theorem 1.4.3]), the operator  $A$  is dissipative in  $V'$  with  $(Ax|x)_{V'} \leq -\mu|x|_{V'}^2$ , and the function  $f$  defined in Proposition 4.15 can be rewritten by means of (3.6) (5.12) as

$$\begin{aligned}
f(x) &= -B\left(\frac{1}{4\beta_0}(\Psi'(x)[0] - q_0), \frac{1}{4\beta_1(\cdot)}(\Psi'(x) - q_1)\right) \\
&= -\frac{\langle\delta_0, \Psi'(x)\rangle - q_0}{4\beta_0}\delta_0 - \frac{1}{4\beta_1(\cdot)}(\Psi'(x) - q_1).
\end{aligned}$$

Hence, if  $\bar{x}$  is a CLE-equilibrium point, for all  $x \in V'$ , we have

$$(f(x) - f(\bar{x})|x - \bar{x})_{V'} = -\frac{\langle\delta_0, \Psi'(x) - \Psi'(\bar{x})\rangle}{4\beta_0}(\delta_0|x - \bar{x})_{V'} - \left(\frac{1}{4\beta_1(\cdot)}(\Psi'(x) - \Psi'(\bar{x}))\right)|x - \bar{x}\Big|_{V'}.$$

Note that the second term in the above inequality satisfies

$$\left(\frac{1}{4\beta_1(\cdot)}(\Psi'(x) - \Psi'(\bar{x}))\right)|x - \bar{x}\Big|_{V'} \geq \frac{1}{4|\beta_1|_{L^\infty(0, \bar{s})}}(\Psi'(x) - \Psi'(\bar{x})|x - \bar{x})_{V'} \geq 0$$

since  $\beta_1 \in L^\infty(0, \bar{s})$ , and  $\Psi'$  is a monotone operator (as a consequence of the convexity of  $\Psi$ ). That in particular implies

$$(C.1) \quad (f(x) - f(\bar{x})|x - \bar{x})_{V'} \leq -\frac{\langle\delta_0, \Psi'(x) - \Psi'(\bar{x})\rangle}{4\beta_0}(\delta_0|x - \bar{x})_{V'} \leq \frac{[\Psi]_L|\delta_0|_{V'}^2}{4\beta_0}|x - \bar{x}|_{V'}^2,$$

so that (4.18) is satisfied with  $\xi = -\mu + [\Psi]_L|\delta_0|_{V'}^2/(4\beta_0)$ .  $\square$

## REFERENCES

- [1] P. Acquistapace, F. Flandoli, and B. Terreni, *Initial boundary value problems and optimal control for nonautonomous parabolic systems*, SIAM J. Control Optimiz, 29 (1991), 89–118.
- [2] P. Acquistapace, B. Terreni, *Infinite horizon LQR problems for nonautonomous parabolic systems with boundary control*, SIAM J. Control Optimiz, 34 (1996), 1–30.
- [3] P. Acquistapace, B. Terreni, *Classical solutions of nonautonomous Riccati equations arising in parabolic boundary control problems*, Appl. Math. Optim., 39 (1999), 361–409.
- [4] P. Acquistapace, B. Terreni, *Classical solutions of nonautonomous Riccati equations arising in Parabolic Boundary Control problems II*, Appl. Math. Optim., 41 (2000), 199–226.
- [5] Asea P, Zak P (1999) Time-to-build and cycles. Journal of economic dynamics and control 23(8):1155-1175.
- [6] Bambi M (2008) Endogenous growth and time-to-build: The AK case. Journal of Economic Dynamics and Control 32(4):1015-1040.
- [7] Bambi M., Fabbri, G., Gozzi, F., (2012) Optimal policy and consumption smoothing effects in the time-to-build AK model. ECONOMIC THEORY, 50 (3), p. 635-669.
- [8] Barbu, V., *Boundary control problems with convex cost criterion*, SIAMJ. Control and Optim., 18(1980), 227-254.
- [9] V. Barbu, G. Da Prato, “Hamilton–Jacobi Equations in Hilbert Spaces,” Pitman, London, 1983.
- [10] V. Barbu, G. Da Prato, *Hamilton-Jacobi equations in Hilbert spaces; variational and semigroup approach*, Ann. Mat. Pura Appl., IV, 42 (1985), 303-349.
- [11] V. Barbu, G. Da Prato, *A note on a Hamilton-Jacobi equation in Hilbert space*, Nonlinear Anal., 9 (1985), 1337–1345.
- [12] V. Barbu, G. Da Prato and C. Popa, *Existence and uniqueness of the dynamic programming equation in Hilbert spaces*, Nonlinear Anal., n. 3, 7 (1983), 283-299.
- [13] V. Barbu, Th. Precupanu, “Convexity and Optimization in Banach Spaces,” Editura Academiei, Bucharest, 1986.
- [14] E. Barucci, F. Gozzi, *Investment in a Vintage Capital Model*, Research in Economics, Vol 52, pp.159–188, 1998.
- [15] E. Barucci, F. Gozzi, *Technology Adpotion and Accumulation in a Vintage Capital Model*, J. of Economics, Vol. 74, no. 1, pp.1–30, 2001.
- [16] Benhabib, J., Rustichini, A., 1991, Vintage capital, investment, and growth, Journal of Economic Theory, 55, 323-339.
- [17] A. Bensoussan, G. Da Prato, M.C. Delfour, S.K. Mitter, “Representation and Control of Infinite Dimensional Systems,” Vol. 1 & 2, Birkäuser, Boston, 1993.
- [18] Boucekkine, R., Camacho, C., and Fabbri, G. (2013). Spatial dynamics and convergence: The spatial AK model. Journal of Economic Theory, 148(6), 2719-2736.
- [19] Boucekkine, R., Fabbri, G., Federico, S., Gozzi, F., (2018). Growth and Agglomeration in the Heterogeneous Space: A Generalized AK Approach. 2017. [jhalshs-01399995v2](https://halshs.archives-ouvertes.fr/halshs-01399995v2).
- [20] Boucekkine, R., Licandro, O., Christopher, P., 1997, Differential-difference equations in economics: on the numerical solution of vintage capital growth models, Journal of Economic Dynamics and Control, 21, 347-362.
- [21] Boucekkine, R., Germain, M., Licandro, O., Magnus, A., 1998, Creative destruction, investment volatility, and the average age of capital, Journal of Economic Growth, 3 , 361-384.
- [22] Boucekkine, R., del Rio, F., Licandro, O., 1999, Endogenous vs. exogenously driven fluctuations in vintage capital models, Journal of Economic Theory, 88, 161-187.
- [23] Boucekkine, R., Germain, M, Licandro, O., Magnus, A., 2001, Numerical solution by iterative methods of a class of vintage capital models, Journal of Economic Dynamics and Control, 25, 655-669.

- [24] P. Cannarsa, G. Di Blasio, *A direct approach to infinite dimensional Hamilton–Jacobi equations and applications to convex control with state constraints*, *Diff. Int. Eq.*, 8 (1995), no. 2, 225–246.
- [25] P. Cannarsa, F. Gozzi, H.M. Soner, *A dynamic programming approach to nonlinear boundary control problems of parabolic type*. *J. Funct. Anal.* 117 (1993), no. 1, 25–61.
- [26] P. Cannarsa, M.E. Tessitore, *Infinite dimensional Hamilton–Jacobi equations and Dirichlet boundary control problems of parabolic type*, *SIAM J. Control Optim.*, 34 (1996), 1831–1847.
- [27] CannarsaTessitore94
- [28] M.G. Crandall, P.L. Lions, *Hamilton–Jacobi equations in infinite dimensions. Part I: Uniqueness of viscosity solutions*, *J. Funct. Anal.*, 62 (1985), 379–396;  
*Part II: Existence of viscosity solutions*, *J. Funct. Anal.*, 65 (1986), 368–405;  
*Part III*, *J. Funct. Anal.*, 68 (1986), 214–247;  
*Part IV: Hamiltonians with unbounded linear terms*, *J. Funct. Anal.*, 90 (1990), 237–283;  
*Part V: Unbounded linear terms and B–continuous solutions*, *J. Funct. Anal.*, 97, 2, (1991), 417–465;  
*Part VI: Nonlinear A and Tataru’s method refined*, *Evolution equations, control theory, and biomathematics*, *Han sur Lesse*, (1991), 51–89, *Lecture Notes in Pure and Appl. Math.*, 155, Dekker, New York, 1994;  
*Part VII: The HJB equation is not always satisfied*, *J. Func. Anal.*, 125 (1994), 111–148.
- [29] Davidson, R., Harris, R., 1981, *Non-convexities in continuous-time investment theory*, *Review of Economic Studies*, 48. 235–253.
- [30] G. Di Blasio, *Global solutions for a class of Hamilton–Jacobi equations in Hilbert spaces*, *Numer. Funct. Anal. Optim.*, 8 (1985/86), no. 3–4, 261–300.
- [31] G. Di Blasio, *Optimal control with infinite horizon for distributed parameter systems with constrained controls*, *SIAM J. Control & Optim.*, 29 (1991); no. 4, 909–925.
- [32] Eisner, R., Strotz, R.H., 1963, *Determinants of business investment*, in *Impacts of Monetary Policy*, Prentice Hall, Englewood Cliffs, USA, 60–223.
- [33] Engel, K.; Nagel, R., *One-parameter semigroups for linear evolution equations* *Graduate Texts in Mathematics*, 194, Springer-Verlag, New York, 2000.
- [34] Fabbri, G. (2016). *Geographical structure and convergence: A note on geometry in spatial growth models*. *Journal of Economic Theory*, 162(1), 114–136.
- [35] Fabbri, G., Gozzi, F., *Solving optimal growth models with vintage capital: The dynamic programming approach*, *J. Econ. Theory* (2008), vol. 143 , 1, pp. 331–373.
- [36] Fabbri, G., Gozzi, F., Swiech, A., *Stochastic Optimal Control in Infinite Dimension*, Springer, 2017.
- [37] G. Fabbri; S. Faggian; G. Freni, *On the Mitra–Wan Forest Management Problem in Continuous Time*, *J. Econ. Theory* (2015), vol. 157, pp. 1001–1040
- [38] S. Faggian, *Boudary control problems with convex cost and Dynamic Programming in infinite dimension. Part 1: the maximum principle*, *Diff. and Int. Eq.*, Vol. 17 (2004), n.9–10, pp. 1149–1174.
- [39] S. Faggian, *Boudary control problems with convex cost and Dynamic Programming in infinite dimension. Part 2: the HJB equation*, *Discrete Contin. Dyn. Syst.*, 12 (2005), n.2 pp. 323–346.
- [40] S. Faggian, *Regular solutions of Hamilton–Jacobi equations arising in Economics*, *Appl. Math. Optim.*, Vol. 51, (2005), n. 2, pp. 123–162.
- [41] S. Faggian, *Applications of dynamic programming to economic problems with vintage capital*, *Dynamics of Continuous, Discrete and Impulsive Systems Series A: Mathematical Analysis* (2008), 15 (4).
- [42] S. Faggian, *Hamilton–Jacobi equations arising from boundary control problems with state constraints*, *SIAM J Control Optim.*, vol. 47, pp. 2157–217
- [43] S. Faggian, F. Gozzi *On the dynamic programming approach for optimal control problems of PDE’s with age structure*, *Math. Pop Stud.*, Vol 11, n. 3–4, 2004, pp.233–270.

- [44] S. Faggian, F. Gozzi, *Optimal investment models with vintage capital: a Dynamic programming approach*, J. Math. Econ. (2010).
- [45] Fattorini, H. O., *Boundary control systems*. SIAM J. Control, 6, 1968, pp. 349385.
- [46] , Fattorini87
- [47] G. Feichtinger, R. Hartl, and S. Sethi. *Dynamical Optimal Control Models in Advertising: Recent Developments*. Management Sci., 40:195226, 1994.
- [48] G. Feichtinger, G. Tragler, and V.M. Veliov, *Optimality Conditions for Age-Structured Control Systems*, J. Math. Anal. Appl. 288(1), 47-68, 2003.
- [49] G. Feichtinger, R.F. Hartl, P.M. Kort, and V.M. Veliov, *Dynamic investment behavior taking into account ageing of the capital goods*, Dynamical systems and control, 379–391, Stability Control Theory Methods Appl., 22, Chapman & Hall/CRC, Boca Raton, FL, 2004.
- [50] G. Feichtinger, R.F. Hartl, P.M. Kort, and V.M. Veliov. *Anticipation Effects of Technological Progress on Capital Accumulation: a Vintage Capital Approach*, J. Econom. Theory 126 (2006), no. 1, 143–164.
- [51] F. Gozzi, *Some results for an optimal control problem with a semilinear state equation I*, Rendiconti dell'Accademia Nazionale dei Lincei, n. 8, 82 (1988), 423-429.
- [52] F. Gozzi, *Some results for an optimal control problem with a semilinear state equation II*, Siam J. Control and Optim., n. 4, 29 (1991), 751-768.
- [53] F. Gozzi, *Some results for an infinite horizon control problem governed by a semilinear state equation*, Proceedings Vorau, July 10-16 1988; editors F.Kappel, K.Kunisch, W.Schappacher; International Series of Numerical Mathematics, Vol. 91, Birkäuser - Verlag, Basel, 1989, 145-163.
- [54] F. Gozzi, A. Swiech, X.Y. Zhou, *A corrected proof of the stochastic verification theorem within the framework of viscosity solutions*, Siam J. Control and Optim., 43 (2005), no. 6, 2009–2019.
- [55] Gozzi, F.; Tessitore, M. E., *Sufficient conditions for Dirichlet boundary control problems of parabolic type*, Partial differential equation methods in control and shape analysis (Pisa), 189204, Lecture Notes in Pure and Appl. Math., 188, Dekker, New York, 1997.
- [56] . Gozzi, F.; Tessitore, M. E., *Optimality conditions for Dirichlet boundary control problems of parabolic type* J. Math. Systems Estim. Control 8 (1998), no. 1.
- [57] Greenwood, J., Jovanovic, B., 2001, Accounting for growth, Studies in Income and Wealth: New Directions in Productivity Analysis, University of Chicago Press, Chicago, USA.
- [58] Iftode, V. *Hamilton-Jacobi equations an dboundar yconvex control problems*, Rev. Roumain eMath. Pures Appl., 34(2), (1989), 117-127.
- [59] Kato, N., A principle of linearized stability for nonlinear evolution equations, Transactions of the American Mathematical society, 347 (1995), pp.2851-2868.
- [60] I. Lasiecka, R. Triggiani, “Control Theory for Partial Differential Equations: Continuous and Approximation Theory,” Vol I. Abstract parabolic systems. Encyclopedia of Mathematics and its Applications, 74. Vol II. Abstract hyperbolic-like systems over a finite time horizon; Encyclopedia of Mathematics and its Applications; CambridgeUniversity Press: Cambridge, 2000, 74-75.
- [61] I. Lasiecka and R. Triggiani. Differential and algebraic equations with application to boundary control problems: Continuous theory and approximation theory, Lecture Notes in Control and Information Sciences. New York: Springer-Verlag, 1991,p. 164.
- [62] Lions, J.-L., Optimal control of systems governed by partial differential equations. Translated from the French by S. K. Mitter. Die Grundlehren der mathematischen Wissenschaften, Band 170 Springer-Verlag, New York-Berlin 1971.
- [63] X.Y. Li, J.M. Yong *Optimal Control Theory for Infinite Dimensional Systems*, Birkhauser, Boston, Cambridge, MA, 1995.

- [64] Lunardi, A., *Analytic Semigroups, Optimal Regularity and Parabolic Problems*, Birkhauser, Boston, Cambridge, MA, 1995.
- [65] Malcomson, J.M., 1975, Replacement and the rental value of capital equipment subject to obsolescence, *Journal of Economic Theory*, 10, 24-41.
- [66] C. Marinelli, *Optimal advertising under uncertainty*, PHD thesis, Columbia University, 2003.
- [67] Moll, B., Nuno, G., Social optima in economies with heterogeneous agents, *Review of Economic Dynamics* 28 (2018), 150180.
- [68] Pazy, A. *Semigroups of linear operators and applications to partial differential equations*. Applied Mathematical Sciences, 44. Springer-Verlag, New York, 1983.
- [69] Solow, R., Tobin, J., Von Weizsacker, C.C., Yaari, M., 1966, Neoclassical growth with fixed factor productions, *Review of Economic Studies*, 33, 79-115.
- [70] Tröltzsch, F., *On the semigroup approach for the optimal control of semilinear parabolic equations including distributed and boundary control*. *Z. Anal. Anwendungen* 8 (1989), no. 5, 431443.
- [71] Tröltzsch, F., Tröltzsch, Fredi *Optimal control of partial differential equations. Theory, methods and applications*. Translated from the 2005 German original by Jrgen Sprekels. Graduate Studies in Mathematics, 112. American Mathematical Society, Providence, RI, 2010.
- [72] Xepapadeas, A. De Zeeuw, A., 1999, Environmental policy and competitiveness: the Porter Hypothesis and the composition of capital, *Journal of Environmental Economics and Management*, 37,165-182.
- [73] J. Yong, X.Y. Zhou, “Stochastic controls. Hamiltonian systems and HJB equations. Applications of Mathematics,” 43. Springer-Verlag, New York, 1999.

[1]S. FAGGIAN, DEPARTMENT OF ECONOMICS, UNIVERSITÁ “Ca’ Foscari” VENEZIA, ITALY. FAGGIAN@UNIVE.IT.

[2] DIPARTIMENTO DI ECONOMIA E FINANZA, UNIVERSITÁ LUISS - “GUIDO CARLI, I-00162, ROMA, ITALY. FGOZZI@LUISS.IT.

[3] CENTER, DEPARTMENT OF ECONOMETRICS & OPERATIONS RESEARCH, TILBURG UNIVERSITY, P.O. BOX 90153, 5000 LE TILBURG, THE NETHERLANDS; DEPARTMENT OF ECONOMICS, UNIVERSITY OF ANTWERP, PRINSSTRAAT 13, 2000 ANTWERP 1, BELGIUM, KORT@UVT.NL.